

Exact solutions of nonlinear partial differential equations by singularity analysis

R. Conte

Service de physique de l'état condensé, CEA Saclay,
F-91191 Gif-sur-Yvette Cedex, France

Abstract. Whether integrable, partially integrable or nonintegrable, nonlinear partial differential equations (PDEs) can be handled from scratch with essentially the same toolbox, when one looks for analytic solutions in closed form. The basic tool is the appropriate use of the singularities of the solutions, and this can be done without knowing these solutions in advance. Since the elaboration of the *singular manifold method* by Weiss *et al.*, many improvements have been made. After some basic recalls, we give an interpretation of the method allowing us to understand why and how it works. Next, we present the state of the art of this powerful technique, trying as much as possible to make it a (computerizable) algorithm. Finally, we apply it to various PDEs in $1 + 1$ dimensions, mostly taken from physics, some of them chaotic : sine-Gordon, Boussinesq, Sawada-Kotera, Kaup-Kupershmidt, complex Ginzburg-Landau, Kuramoto-Sivashinsky, etc.

PACS : 02.90.+p, 03.40.Kf

Keywords

Exact solutions
Nonperturbative methods
Painlevé analysis

Direct and inverse methods in nonlinear evolution equations, CIME school,
Cetraro, 5–12 September 1999, ed. A. Greco, Springer Verlag (2000).

19 February 2000

S99/072, nlin-si/0009024

1 Introduction

Our interest is to find explicitly the “macroscopic” quantities which materialize the integrability of a given *nonlinear* differential equation, such as particular solutions or first integrals. We mainly handle partial differential equations (PDEs), although some examples are taken from ordinary differential equations (ODEs). Indeed, the methods described in these lectures apply equally to both cases.

These methods are based on the *a priori* study of the singularities of the solutions. The reader is assumed to possess a basic knowledge of the singularities of nonlinear *ordinary* differential equations, the Painlevé property for ODEs and the Painlevé test. All this prerequisite material is well presented in a book by Hille [57] while Cargèse lecture notes [26] contain a detailed exposition of the methods, including the Painlevé test for ODEs. Many applications are given in a review [102].

As a general bibliography on the subject of these lectures, we recommend Cargèse lecture notes [84] and a shorter subset of these with emphasis on the various so-called truncations [24].

Throughout the text, we exclude linear equations, unless explicitly stated.

2 Various levels of integrability for PDEs, definitions

In this section, we review the required definitions (exact solution, Bäcklund transformation, Lax pair, Darboux transformation, etc).

The most important point is the global nature of the information which is looked for. The existence theorem of Cauchy (for ODEs) or Cauchy-Kowalevski (for PDEs) is of no help for this purpose. Indeed, it only states a local property and says nothing on what happens outside the disk of definition of the Taylor series. Therefore it cannot distinguish between chaotic equations and integrable ones.

Still from this point of view, Laurent series are not better than Taylor series. For instance, the Bianchi IX cosmological model is a six-dimensional dynamical system

$$\sigma^2(\text{Log } A)'' = A^2 - (B - C)^2, \quad \text{and cyclically, } \sigma^4 = 1, \quad (1)$$

which is undoubtedly chaotic [107]. Despite the existence of the Laurent series [39]

$$\begin{aligned} A/\sigma &= \chi^{-1} + a_2\chi + O(\chi^3), \quad \chi = \tau - \tau_1, \\ B/\sigma &= b_0\chi + b_1\chi^2 + O(\chi^3), \\ C/\sigma &= c_0\chi + c_1\chi^2 + O(\chi^3), \end{aligned} \quad (2)$$

which depends on six independent arbitrary coefficients, $(\tau_1, b_0, c_0, b_1, c_1, a_2)$, a wrong statement would be to conclude to the absence of chaos.

This leads us to the definition of the first one of several needed global mathematical objects.

Definition 1. One calls **exact solution** of a nonlinear PDE any solution defined in the whole domain of definition of the PDE and which is given in *closed form*, i.e. as a finite expression.

The opposite of an exact solution is of course not a wrong solution, but what Painlevé calls “une solution illusoire”, such as the above mentioned series.

Note that a multivalued expression is not excluded. From this definition, an exact solution is *global*, as opposed to *local*. This generically excludes series or infinite products, unless their domain of validity can be made the full domain of definition, like for linear ODEs.

Example 2. The Kuramoto-Sivashinsky (KS) equation

$$u_t + uu_x + \mu u_{xx} + \nu u_{xxxx} = 0, \quad \nu \neq 0, \quad (3)$$

describes, for instance, the fluctuation of the position of a flame front, or the motion of a fluid going down a vertical wall, or a spatially uniform oscillating chemical reaction in a homogeneous medium (see Ref. [78] for a review), and it is well known for its chaotic behaviour. An exact solution is the solitary wave of Kuramoto and Tsuzuki [69] in which the wavevector k is fixed

$$u = 120\nu \left(\frac{k}{2} \tanh \frac{k}{2} \xi \right)^3 + \left(\frac{60}{19} \mu - 30\nu k^2 \right) \frac{k}{2} \tanh \frac{k}{2} \xi + c, \\ \xi = x - ct - x_0, \quad k^2 = \frac{11\mu}{19\nu} \text{ or } -\frac{\mu}{19\nu}, \quad (4)$$

which depends on two arbitrary constants (c, x_0) . On the contrary, the Laurent series

$$u = 120\nu \xi^{-3} + \frac{60}{19} \mu \xi^{-1} + c - \frac{120.11}{19^2} \mu^2 \xi + u_6 \xi^3 + O(\xi^4), \quad (5)$$

which depends on three arbitrary constants (c, x_0, u_6) , is not an exact solution, since no closed form expression is yet known for the sum of this series.

There exists a powerful tool to build exact solutions, this is the Bäcklund transformation. For simplicity, but this is not a restriction, we give the basic definitions for a PDE defined as a single scalar equation for one dependent variable u and two independent variables (x, t) .

Definition 3. (Refs. [41] vol. III chap. XII, [80]) A **Bäcklund transformation** (BT) between two given PDEs

$$E_1(u, x, t) = 0, \quad E_2(U, X, T) = 0 \quad (6)$$

is a pair of relations

$$F_j(u, x, t, U, X, T) = 0, \quad j = 1, 2 \quad (7)$$

with some transformation between (x, t) and (X, T) , in which F_j depends on the derivatives of $u(x, t)$ and $U(X, T)$, such that the elimination of u (resp. U) between (F_1, F_2) implies $E_2(U, X, T) = 0$ (resp. $E_1(u, x, t) = 0$). The BT is called the **auto-BT** or the **hetero-BT** according as the two PDEs are the same or not.

Example 4. The sine-Gordon equation (we identify sine-Gordon and sinh-Gordon since an affine transformation on u does not change the integrability nor the singularity structure)

$$\text{sine-Gordon} : E(u) \equiv u_{xt} + 2a \sinh u = 0 \quad (8)$$

admits the auto-BT

$$(u + U)_x + 4\lambda \sinh \frac{u - U}{2} = 0, \quad (9)$$

$$(u - U)_t - \frac{2a}{\lambda} \sinh \frac{u + U}{2} = 0, \quad (10)$$

in which λ is an arbitrary complex constant, called the *Bäcklund parameter*.

Given the obvious solution $U = 0$ (called *vacuum*), the two equations (9)–(10) are Riccati ODEs with constant coefficients for the unknown $e^{u/2}$,

$$(e^{u/2})_x = \lambda(1 - (e^{u/2})^2), \quad (11)$$

$$(e^{u/2})_t = -a(1 - (e^{u/2})^2)/(2\lambda), \quad (12)$$

therefore their general solution is known in closed form

$$e^{u/2} = \tanh \theta, \quad \theta = \left(\lambda x - \frac{a}{2\lambda} t - z_0 \right), \quad (13)$$

with (λ, z_0) arbitrary. This solution is called the *one-soliton solution*, it is also written as

$$\tanh(u/4) = -e^{-2\theta}, \quad u_x = 4\lambda \operatorname{sech} 2\theta, \quad u_t = -2a\lambda^{-1} \operatorname{sech} 2\theta. \quad (14)$$

By iteration, this procedure gives rise to the N -soliton solution [70,1], an exact solution depending on $2N$ arbitrary complex constants (N values of the Bäcklund parameter λ , N values of the shift z_0), with N an arbitrary positive integer. A remarkable feature of the SG-equation, due to the fact that at least one of the two ODEs (9)–(10) is of order one, is that this N -soliton can be obtained from N different copies of the one-soliton by a simple algebraic operation, i.e. without integration (see Musette's lecture [85]).

Example 5. The Liouville equation

$$\text{Liouville: } E(u) \equiv u_{xt} + \alpha e^u = 0 \tag{15}$$

admits two BTs. The first one

$$(u - v)_x = \alpha \lambda e^{(u+v)/2}, \tag{16}$$

$$(u + v)_t = -2\lambda^{-1} e^{(u-v)/2}, \tag{17}$$

is a BT to a linearizable equation called the d'Alembert equation

$$\text{d'Alembert: } E(v) \equiv v_{xt} = 0. \tag{18}$$

The second one is an auto-BT

$$(u + U)_x = -4\lambda \sinh \frac{u - U}{2}, \tag{19}$$

$$(u - U)_t = \lambda^{-1} \alpha e^{(u+U)/2}. \tag{20}$$

The first of these two BTs allows one to obtain the general solution of the nonlinear Liouville equation, see Section 7.

This ideal situation (generation of the general solution) is exceptional and the generic case is the generation of particular solutions only, as in the sine-Gordon example.

The importance of the BT is such that it is often taken as a definition of *integrability*.

Definition 6. A PDE in N independent variables is **integrable** if at least one of the following properties holds.

1. It is linearizable.
2. For $N > 1$, it possesses an auto-BT which, if $N = 2$, depends on an arbitrary complex constant, the Bäcklund parameter.
3. It possesses a hetero-BT to another integrable PDE.

Although partially integrable and nonintegrable equations, i.e. the majority of physical equations, admit no BT, they retain part of the properties of (fully) integrable PDEs, and this is why the methods presented in these lectures apply to both cases as well. For instance, the KS equation admits the *vacuum* solution $u = 0$ and, in Section 2, an iteration will be built leading from $u = 0$ to the solitary wave (4); the nonintegrability manifests itself in the finite number of times this iteration provides a new result ($N = 1$ for the KS equation, and one cannot go beyond (4) [29]).

For various applications of the BT, see Ref. [104].

When a PDE has some good reasons to possess such features, such as the reasons developed in Section 4, we want to find the BT if it exists, since this is a generator of exact solutions, or a degenerate form of the BT if the BT does not exist, and we want to do it by singularity analysis *only*.

Before proceeding, we need to define some other elements of integrability.

Definition 7. Given a PDE, a **Lax pair** is a system of two linear differential operators

$$\text{Lax pair : } L_1(U, \lambda), L_2(U, \lambda), \quad (21)$$

depending on a solution U of the PDE and, in the $1 + 1$ -dimensional case, on an arbitrary constant λ , called the **spectral parameter**, with the property that the vanishing of the commutator $[L_1, L_2]$ is equivalent to the vanishing of the PDE $E(U) = 0$.

A Lax pair can be represented in several, equivalent ways.

The *Lax representation* [72] is a pair of linear operators (L, P) (scalar or matrix) defined by

$$L_1 = L - \lambda, L_2 = \partial_t - P, L_1\psi = 0, L_2\psi = 0, \lambda_t = 0, \quad (22)$$

in which the elimination of the scalar λ yields

$$L_t = [P, L], \quad (23)$$

i.e. , thanks to the *isospectral* condition $\lambda_t = 0$, a time evolution analogous to the one in Hamiltonian dynamics.

The *zero-curvature representation* is a pair (L, M) of linear operators independent of (∂_x, ∂_t)

$$\begin{aligned} L_1 &= \partial_x - L, L_2 = \partial_t - M, L_1\psi = 0, L_2\psi = 0, \\ [\partial_x - L, \partial_t - M] &= L_t - M_x + LM - ML = 0. \end{aligned} \quad (24)$$

The common order N of the matrices is called the *order* of the Lax pair.

The *projective Riccati representation* is a first order system of $2N - 2$ Riccati equations in the unknowns $\psi_j/\psi_1, j = 2, \dots, N$, equivalent to the zero-curvature representation (24).

The *scalar representation* is a pair of scalar linear PDEs, one of them of order higher than one,

$$\begin{aligned} L_1\psi &= 0, L_2\psi = 0, \\ X &\equiv [L_1, L_2] = 0. \end{aligned} \quad (25)$$

In $1 + 1$ -dimensions, one of the PDEs can be made an ODE (i.e. involving only x - or t -derivatives), in which case the order of this ODE is called the order of the Lax pair.

The *string representation* or *Sato representation* [64]

$$[P, Q] = 1. \quad (26)$$

This very elegant representation, reminiscent of Hamiltonian dynamics, uses the Sato definition of a *microdifferential operator* (a differential operator

with positive and negative powers of the differential operator ∂) and of its *differential part* denoted $(\)_+$ (the subset of its nonnegative powers), e.g.

$$Q = \partial_x^2 - u, \quad (27)$$

$$L = Q^{1/2}, \quad (28)$$

$$(L^3)_+ = \partial_x^3 - (3/4)\{u, \partial_x\}, \quad (29)$$

$$(L^5)_+ = \partial_x^5 - (5/4)\{u, \partial_x^3\} + (5/16)\{3u^2 + u_{xx}, \partial_x\}, \quad (30)$$

in which $\{a, b\}$ denotes the anticommutator $ab + ba$. See Ref. [42] for a tutorial presentation.

Example 8. The sine-Gordon equation (8) admits the zero-curvature representation

$$(\partial_x - L) \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = 0, \quad L = \begin{pmatrix} U_x/2 & \lambda \\ \lambda & -U_x/2 \end{pmatrix}, \quad (31)$$

$$(\partial_t - M) \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = 0, \quad M = -(a/2)\lambda^{-1} \begin{pmatrix} 0 & e^U \\ e^{-U} & 0 \end{pmatrix}, \quad (32)$$

equivalent to the Riccati representation, with $y = \psi_1/\psi_2$,

$$y_x = \lambda + U_x y - \lambda y^2, \quad (33)$$

$$y_t = -\frac{a}{2}\lambda^{-1}e^U + \frac{a}{2}\lambda^{-1}e^{-U}y^2. \quad (34)$$

Example 9. The matrix nonlinear Schrödinger equation

$$iQ_t + (b/a)Q_{xx} - 2abQRQ = 0, \quad -iR_t + (b/a)R_{xx} - 2abRQR = 0, \quad (35)$$

in which (Q, R) are rectangular matrices of respective orders (m, n) and (n, m) , and (i, a, b) constants, admits the zero-curvature representation ([77] Eq. (5))

$$(\partial_x - L)\psi = 0, \quad (\partial_t - M)\psi = 0, \quad (36)$$

$$L = aP + \lambda G, \quad M = (-aGP^2 + GP_x + 2\lambda P + (2/a)\lambda^2 G)b/i, \quad (37)$$

in which λ is the spectral parameter, P and G matrices of order $m+n$ defined as

$$P = \begin{pmatrix} 0 & Q \\ -R & 0 \end{pmatrix}, \quad G = \begin{pmatrix} 1_m & 0 \\ 0 & -1_n \end{pmatrix}. \quad (38)$$

The matrix G characterizes the internal symmetry group $\text{GL}(m, \mathcal{C}) \otimes \text{GL}(n, \mathcal{C})$. The lowest values

$$m = 1, n = 1, \quad Q = (u), \quad R = (U), \quad (39)$$

define the AKNS system (Section 9.1), whose reduction $U = \bar{u}$ is the usual scalar nonlinear Schrödinger equation.

Example 10. The 2 + 1-dimensional Ito equation [62]

$$E(u) \equiv (u_{xxxx} + 6\alpha^{-1}u_{xt}u_{xx} + a_1u_{tt} + a_2u_{xt} + a_3u_{xx} + a_4u_{ty})_x = 0 \quad (40)$$

has a Lax pair whose scalar representation is

$$L_1 \equiv \partial_x^3 + a_1\partial_t + (a_2 + 6\alpha^{-1}U_{xx})\partial_x + a_4\partial_y - \lambda \quad (41)$$

$$L_2 \equiv \partial_x\partial_t - \mu\partial_x + \left(\frac{a_3}{3} + 2\alpha^{-1}U_{xt}\right) \quad (42)$$

$$\alpha[L_1, L_2] = 2E(U) + 6U_{xxx}L_2. \quad (43)$$

In the 2 + 1-dimensional case $a_4 \neq 0$, the parameter λ can be set to 0 by the change $\psi \mapsto \psi e^{\lambda y}$. This is the reason of the precision at the end of item 2 in definition 7. This pair has the order four in the generic case $a_1 \neq 0$, although neither L_1 nor L_2 has such an order.

Example 11. The string representation of the Lax pair of the derivative of the first Painlevé equation is

$$[P, Q] = [((\partial_x^2 - u)^3)_+, \partial_x^2 - u] = -(1/4)u_{xxx} + (3/4)uu_x = 1. \quad (44)$$

Example 12. The Sato representation of the Lax pair for the whole Korteweg-de Vries hierarchy is

$$\partial_{t_m}L = [(L^{2m-1})_+, L], \quad L = Q^{1/2}, \quad Q = \partial_x^2 - u, \quad m = 1, 3, 5, \dots \quad (45)$$

From the singularity point of view, the Riccati representation is the most suitable, as will be seen.

The last main definition we need is the Darboux transformation (DT). The working definition given below is very simplified (this is an involution) as compared to the one of Darboux [40], but it is sufficient for our purpose. The full definition is given in Musette's lecture [85].

Definition 13. Given a PDE, a **Darboux transformation** is a transformation between two solutions (u, U) of the PDE

$$\text{DT} : u = \sum_f \mathcal{D}_f \text{Log } \tau_f + U \quad (46)$$

linking their difference to a finite number of linear differential operators \mathcal{D}_f (f like family) acting on the logarithm of functions τ_f .

In the definition (46), it is important to note that, despite the notation, each function τ_f is in fact the ratio of the “tau-function” of u by that of U .

Lax pairs, Bäcklund and Darboux transformations are not independent. In order to exhibit their interrelation, one needs an additional information, namely the link

$$\forall f : \mathcal{D}_f \text{Log } \tau_f = F_f(\psi), \quad (47)$$

which most often is the identity $\tau = \psi$, between the functions τ_f and the function ψ in the definition of a scalar Lax pair.

Example 14. The (integrable) sine-Gordon equation (8) admits the Darboux transformation

$$u = U - 2(\text{Log } \tau_1 - \text{Log } \tau_2), \quad (48)$$

in which (τ_1, τ_2) is a solution (ψ_1, ψ_2) of the system (31)–(32).

Then its BT (9)–(10) is the result of the elimination [14] of τ_1/τ_2 between the DT (48) and the Riccati form of the Lax pair (33)–(34), with the correspondence $\tau_f = \psi_f, f = 1, 2$. This elimination reduces to the substitution $y = e^{-(u-U)/2}$ in the Riccati system (33)–(34), and this is one of the advantages of the Riccati representation. Therefore the Bäcklund parameter and the spectral parameter are identical notions.

Example 15. The (nonintegrable) Kuramoto-Sivashinsky equation admits the degenerate Darboux transformation

$$u = U + (60\nu\partial_x^3 + (60/19)\mu\partial_x)\text{Log } \tau, \quad (49)$$

in which $U = c$ (vacuum) and τ is the general solution ψ of the linear system (a degenerate second order scalar Lax pair)

$$L_1\psi \equiv (\partial_x^2 - k^2/4)\psi = 0, \quad (50)$$

$$L_2\psi \equiv (\partial_t + c\partial_x)\psi = 0, \quad (51)$$

$$[L_1, L_2] \equiv 0. \quad (52)$$

The solution u defined by (49) is then the solitary wave (4), and this is a much simpler way to write it, because the logarithmic derivatives in (49) take account of the whole nonlinearity.

Since, roughly speaking, the BT is equivalent to the couple (DT, Lax pair), one can rephrase as follows the iteration to generate new solutions. Let us symbolically denote

- $E(u) = 0$ the PDE,
- $\text{Lax}(\psi, \lambda, U) = 0$ a scalar Lax pair,
- F the link (47) $\mathcal{D}\text{Log } \tau = F(\psi)$ from ψ to τ ,
- $u = \text{Darboux}(U, \tau)$ the Darboux transformation.

The iteration is the following, see e.g. [54].

1. (initialization) Choose $u_0 =$ a particular solution of $E(u) = 0$; set $n = 1$; perform the following loop until some maximal value of n ;
2. (start of loop) Choose $\lambda_n =$ a particular complex constant;
3. Compute, by integration, a particular solution ψ_n of the linear system $\text{Lax}(\psi, \lambda_n, u_{n-1}) = 0$;
4. Compute, without integration, $\mathcal{D}\text{Log } \tau_n = F(\psi_n)$;
5. Compute, without integration, $u_n = \text{Darboux}(u_{n-1}, \tau_n)$;
6. (end of loop) Set $n = n + 1$.

Depending on the choice of λ_n at step 2, and of ψ_n at step 3, one can generate either the N -soliton solution, or solutions rational in (x, t) , or a mixture of such solutions.

3 Importance of the singularities : a brief survey of the theory of Painlevé

A classical theorem states that a function of one complex variable without any singularity in the analytic plane (i.e. the complex plane compactified by addition of the unique point at infinity) is a constant. Therefore a function with singularities is characterized, as shown by Mittag-Leffler, by the knowledge of its singularities in the analytic plane. Similarly, if u satisfies an ODE or a PDE, the structure of singularities of the general solution characterizes the level of integrability of the equation. This is the basis of the theory of the (explicit) integration of nonlinear ODEs built by Painlevé, which we only briefly introduce here [for a detailed introduction, see Cargèse lecture notes : Ref. [26] for ODEs, Ref. [84] for PDEs].

To integrate an ODE is to acquire a global knowledge of its general solution, not only the local knowledge ensured by the existence theorem of Cauchy. So, the most demanding possible definition for the “integrability” of an ODE is the single valuedness of its general solution, so as to adapt this solution to any kind of initial conditions. Since even linear equations may fail to have this property, e.g. $2xu' + u = 0$, $u = cx^{-1/2}$, a more reasonable definition is the following one.

Definition 16. The *Painlevé property* (PP) of an ODE is the uniformizability of its general solution.

In the above example, the uniformization is achieved by the change of independent variable $x = X^2$. This definition is equivalent to the more familiar one.

Definition 17. The *Painlevé property* (PP) of an ODE is the absence of movable critical singularities in its general solution.

Definition 18. The *Painlevé property* (PP) of a PDE is its integrability (Definition 6) and the absence of movable critical singularities in its general solution.

Let us recall that a singularity is said *movable* (as opposed to *fixed*) if its location depends on the initial conditions, and *critical* if multivaluedness takes place around it. Indeed, out of the four configurations of singularities (critical or noncritical) and (fixed or movable), only the configuration (critical and movable) prevents uniformizability : one does not know where to put the cut since the point is movable.

Wrong definitions of the PP, alas repeatedly published, consist in replacing in the definition “movable critical singularities” by “movable singularities other than poles”, or “its general solution” by “all its solutions”. Even worse definitions only refer to Laurent series. See Ref. [26], Section 2.6, for the arguments of Painlevé himself.

The mathematicians like Painlevé want to integrate whole classes of ODEs (e.g. second order algebraic ODEs). We will only use their methods for a given ODE or PDE, with the aim of deriving the elements of integrability described in Section 2 (exact solutions, ...). This *Painlevé analysis* is twofold (“double méthode”, says Painlevé).

1. Build necessary conditions for an ODE or a PDE to have the PP (this is called the *Painlevé test*).
2. When all these conditions are satisfied, or at least some of them, find the global elements of integrability. In the integrable case this is achieved either (ODE case) by explicitly integrating or (PDE case) by finding an auto-BT (like equations (9)–(10) for sine–Gordon) or a BT towards another PDE with the PP (like (16)–(17) between the d’Alembert and Liouville equations). In the partially integrable case, only degenerate forms of the above can be expected, as described in Section 2.

4 The Painlevé test for PDEs in its invariant version

When the PDE reduces to an ODE, the Painlevé test (for shortness we will simply say the test) reduces by construction to the test for ODEs, presented in detail elsewhere [26] and assumed known here.

We will skip those steps of the test which are the same for ODEs and for PDEs (e.g., diophantine conditions that all the leading powers and all the Fuchs indices be integral), and we will concentrate on the features which are specific to PDEs, namely the description of the movable singularities, the optimal choice of the expansion variable for the Laurent series, the advantage of the homographic invariance.

4.1 Singular manifold variable φ and expansion variable χ

Consider a nonlinear PDE

$$E(u, x, t, \dots) = 0. \quad (53)$$

To test movable singularities for multivaluedness without integrating, which is the essence of the test, one must first describe them, then, among other steps, check the existence near each movable singularity of a Laurent series which represents the general solution.

For PDEs, the singularities are not isolated in the space of the independent variables (x, t, \dots) , but they lay on a codimension one manifold

$$\varphi(x, t, \dots) - \varphi_0 = 0, \quad (54)$$

in which the *singular manifold variable* φ is an arbitrary function of the independent variables and φ_0 an arbitrary movable constant. Even in the

ODE case, the movable singularity can be defined as $\varphi(x) - \varphi_0 = 0$, since the implicit functions theorem allows this to be locally inverted to $x - x_0 = 0$; the arbitrary function φ thus introduced may then be used to construct exact solutions which would be impossible to find with the restriction $\varphi(x) = x$ [114,92].

One must then define from $\varphi - \varphi_0$ an *expansion variable* χ for the Laurent series, for there is no reason to confuse the roles of the singular manifold variable and the expansion variable. Two requirements must be respected: firstly, χ must vanish as $\varphi - \varphi_0$ when $\varphi \rightarrow \varphi_0$; secondly, the structure of singularities in the φ complex plane must be in a one-to-one correspondence with that in the χ complex plane, so χ must be a homographic transform of $\varphi - \varphi_0$ (with coefficients depending on the derivatives of φ).

The Laurent series for u and E involved in the Kowalevski-Gambier part of the test are defined as

$$u = \sum_{j=0}^{+\infty} u_j \chi^{j+p}, \quad -p \in \mathcal{N}, \quad E = \sum_{j=0}^{+\infty} E_j \chi^{j+q}, \quad -q \in \mathcal{N}^* \quad (55)$$

with coefficients u_j, E_j independent of χ and only depending on the derivatives of φ .

To illustrate our point, let us take as an example the Korteweg-de Vries equation

$$E \equiv bu_t + u_{xxx} - (6/a)uu_x = 0 \quad (56)$$

(this is one of the very rare locations where this equation can be taken as an example; indeed, usually, things work so nicely for KdV that it is hazardous to draw general conclusions from its single study).

The choice $\chi = \varphi - \varphi_0$ originally made by Weiss *et al.* [119] makes the coefficients u_j, E_j invariant under the two-parameter group of translations $\varphi \mapsto \varphi + b'$, with b' an arbitrary complex constant and therefore they only depend on the differential invariant $\text{grad } \varphi$ of this group and its derivatives :

$$u = 2a\varphi_x^2\chi^{-2} - 2a\varphi_{xx}\chi^{-1} + ab\frac{\varphi_t}{6\varphi_x} + \frac{2a}{3}\frac{\varphi_{xxx}}{\varphi_x} - \frac{a}{2}\left[\frac{\varphi_{xx}}{\varphi_x}\right]^2 + O(\chi), \quad \chi = \varphi - \varphi_0. \quad (57)$$

There exists a choice of χ for which the coefficients exhibit the highest invariance and therefore are the shortest possible (all details are in Section 6.4 of Ref. [26]), this best choice is [19]

$$\chi = \frac{\varphi - \varphi_0}{\varphi_x - \frac{\varphi_{xx}}{2\varphi_x}(\varphi - \varphi_0)} = \left[\frac{\varphi_x}{\varphi - \varphi_0} - \frac{\varphi_{xx}}{2\varphi_x} \right]^{-1}, \quad \varphi_x \neq 0, \quad (58)$$

in which x denotes one of the independent variables whose component of $\text{grad } \varphi$ does not vanish. The expansion coefficients u_j, E_j are then invariant

under the six-parameter group of homographic transformations

$$\varphi \mapsto \frac{a'\varphi + b'}{c'\varphi + d'}, \quad a'd' - b'c' \neq 0, \quad (59)$$

in which a', b', c', d' are arbitrary complex constants. Accordingly, these coefficients only depend on the following elementary differential invariants and their derivatives: the *Schwarzian*

$$S = \{\varphi; x\} = \frac{\varphi_{xxx}}{\varphi_x} - \frac{3}{2} \left(\frac{\varphi_{xx}}{\varphi_x} \right)^2, \quad (60)$$

and one other invariant per independent variable t, y, \dots

$$C = -\varphi_t/\varphi_x, \quad K = -\varphi_y/\varphi_x, \quad \dots \quad (61)$$

The reason for the minus sign in the definition of C is that, under the travelling wave reduction $\xi = x - ct$, the variable C becomes the constant c . These two invariants are linked by the cross-derivative condition

$$X \equiv ((\varphi_{xxx})_t - (\varphi_t)_{xxx})/\varphi_x = S_t + C_{xxx} + 2C_x S + C S_x = 0, \quad (62)$$

identically satisfied in terms of φ .

For our KdV example, the final Laurent series, as compared with the initial one (57), is remarkably simple :

$$u = 2a\chi^{-2} - ab\frac{C}{6} + \frac{2aS}{3} - 2a(bC - S)_x\chi + O(\chi^2), \quad \chi = (58). \quad (63)$$

For the practical computation of (u_j, E_j) as functions of (S, C) only, i.e. what is called the invariant Painlevé analysis, the above explicit expressions of (S, C, χ) in terms of φ are *not* required, the variable φ completely disappears, and the only necessary information is the gradient of the expansion variable χ defined by Eq. (58). This gradient is a polynomial of degree two in χ (this is a property of homographic transformations), whose coefficients only depend on S, C :

$$\chi_x = 1 + \frac{S}{2}\chi^2, \quad (64)$$

$$\chi_t = -C + C_x\chi - \frac{1}{2}(CS + C_{xx})\chi^2. \quad (65)$$

The above choice (58) of χ which generates homographically invariant coefficients is the simplest one, but it is only particular. The general solution to the above two requirements which also generates homographically invariant coefficients is defined by an affine transformation on the inverse of χ [86]

$$Y^{-1} = B(\chi^{-1} + A), \quad B \neq 0. \quad (66)$$

Since a homography conserves the Riccati nature of an ODE, the variable Y satisfies a Riccati system, easily deduced from the canonical one (64)–(65) satisfied by χ , see (115)–(116).

A frequent worry is : is there any restriction (or advantage, or inconvenient) to perform the test with χ or Y rather than with $\varphi - \varphi_0$? The precise answer is : the three Laurent series are equivalent (their set of coefficients are in a one-to-one correspondence, only their radii of convergence are different). As a consequence, the Painlevé test, which involves the *infinite* series, is insensitive to the choice, and the costless choice (the one which minimizes the computations) is undoubtedly χ defined by its gradient (64)–(65) (to perform the test, one can even set, following Kruskal [63], $S = 0, C_x = 0$). If the same question were asked not about the test but about the second stage of Painlevé analysis as formulated at the end of Section 3, the answer would be quite different, and it is given in Section 6.1.

Finally, let us mention a useful technical simplification. From its definition (58), the variable χ^{-1} is a logarithmic derivative, so the system (64)–(65) can be integrated once

$$\Psi = (\varphi - \varphi_0)\varphi_x^{-1/2}, \quad (67)$$

$$(\text{Log } \Psi)_x = \chi^{-1}, \quad (68)$$

$$(\text{Log } \Psi)_t = -C\chi^{-1} + \frac{1}{2}C_x. \quad (69)$$

This feature helps to process PDEs which can be defined in either conservative or potential form when the conservative field u has a simple pole, such as the Burgers equation

$$E(u) \equiv bu_t + (u^2/a + u_x)_x = 0, \quad (70)$$

$$F(v) \equiv bv_t + v_x^2/a + v_{xx} + G(t) = 0, \quad u = v_x, \quad E = F_x. \quad (71)$$

Despite its (unique) logarithmic term, the ψ -series for v

$$v = a \text{Log } \Psi + v_0 + (2v_{0,x} - abC)\chi + (F(v_0) - aS/2 + abC_x/2)\chi^2 + O(\chi^3), \quad (72)$$

in which v_0 is arbitrary, is “shorter” than the Laurent series for u

$$u = a\chi^{-1} + (ab/2)C + u_2\chi + [(a/4)(b^2(C_t + CC_x) + 2bC_{xx} - S_x - u_{2,x})]\chi^2 + O(\chi^3), \quad (73)$$

in which u_2 is arbitrary, and the resulting series for $F(v)$, which is not a ψ -series but a Laurent series, is *much* shorter than the Laurent series for $E(u)$. See Section 7.3 for an application.

4.2 The WTC part of the Painlevé test for PDEs

As mentioned at the beginning of Section 4, we do not give here all the detailed steps of the test nor all the necessary conditions which it generates (this is done in Section 6.6 of Ref. [26]). We mainly state the notation to be extensively used throughout next sections.

The WTC part [119] of the full test, when rephrased in the equivalent invariant formalism [23], consists in checking the existence of all Laurent series (55) able to represent the general solution, maybe after suitable perturbations [28,89] not describe here.

The gradient of the expansion variable χ is given by (64)–(65), with the cross-derivative condition (62). This condition may be used to eliminate, depending on the PDE, either derivatives $S_{mx,nt}$, with $n \geq 1$, or derivatives $C_{mx,nt}$, with $m \geq 3$, and all equations later written are already simplified in either way.

The *first step* is to find all the admissible values (p, u_0) which define the leading term of the series for u . Such an admissible couple is called a *family of movable singularities* (the term *branch* should be avoided for the confusion which it induces with branching, i.e. multivaluedness).

The recurrence relation for the next coefficients u_j , after replacement of (p, u_0) ,

$$\forall j \geq 1 : E_j \equiv P(j)u_j + Q_j(\{u_l \mid l < j\}) = 0 \tag{74}$$

depends linearly on u_j and nonlinearly on the previously computed coefficients u_l .

The *second step* is to compute the *indicial equation*

$$P(i) = 0 \tag{75}$$

(a determinant in the multidimensional case of a system of PDEs). Its roots are called the *Fuchs indices* of the family because they are indeed the characteristic indices of a linear differential equation near a Fuchsian singularity (the name *resonances* sometimes given to these indices refers to no resonance phenomenon and should also be avoided). One then requires that all indices be integral and obey a rank condition which, for a single PDE, reduces to the condition that all indices be distinct. The value $i = -1$ is always a Fuchs index.

The *third and last step* is to require that, for any admissible family and any Fuchs index i (a signed integer), the *no-logarithm condition*

$$\forall i \in \mathcal{Z}, P(i) = 0 : Q_i = 0 \tag{76}$$

holds true, so that the coefficient u_i is an arbitrary function of the independent variables. In the multidimensional case, this is the condition of orthogonality between the vector \mathbf{Q}_i and the adjoint of the linear operator $\mathbf{P}(i)$. Whenever there exist negative integers in addition to the ever present value

-1 counted with multiplicity one, the condition (76) can only be tested by a perturbation [28].

This ends this subset of the test which, let us insist on the terminology, is only aimed at building *necessary* conditions for the PP.

The Laurent series for u built in this way depends on at most N arbitrary functions (if N denotes the differential order), namely the coefficients u_i introduced at the N Fuchs indices, including φ for the index -1 .

Any item u_j, E_j, Q_j depends, through the elementary invariants (S, C) , on the derivatives of φ up to the order $j+1$, so the dependences are as follows: $u_0 = f(C)$, $u_1 = f(C, C_x, C_t)$, $u_2 = f(C, C_x, C_t, C_{xx}, C_{xt}, C_{tt}, S)$, \dots

Let us take an example.

Example 19. The Kolmogorov-Petrovskii-Piskunov (KPP) equation [67,93]

$$E(u) \equiv bu_t - u_{xx} + 2d^{-2}(u - e_1)(u - e_2)(u - e_3) = 0, \quad e_j \text{ distinct}, \quad (77)$$

encountered in reaction-diffusion systems (an additional convection term uu_x is quite important in physical applications to prey-predator models [105]) possesses the two families (d denotes any square root of d^2)

$$p = -1, \quad u_0 = d, \quad (78)$$

each family has the same two indices $(-1, 4)$, and the Laurent series for each family reads

$$u = d\chi^{-1} + (s_1/3 - (bd/6)C) \quad (79)$$

$$- (d/6) \left((b^2/6)C^2 - 6a_2 - S - bC_x \right) \chi + O(\chi^2), \quad (80)$$

with the notation

$$s_1 = e_1 + e_2 + e_3, \quad a_2 = ((e_2 - e_3)^2 + (e_3 - e_1)^2 + (e_1 - e_2)^2) / (18d^2). \quad (81)$$

At index $i = 4$, the two no-log conditions, one for each sign of d [18],

$$Q_4 \equiv C[(bdC + s_1 - 3e_1)(bdC + s_1 - 3e_2)(bdC + s_1 - 3e_3) - 3b^2d^3(C_t + CC_x)] = 0 \quad (82)$$

are not identically satisfied, so the PDE fails the test.

It is time to define a quantity which, although useless for the test itself, is of first importance at the second stage of Painlevé analysis, which will be developed in Sections 5 to 9. This quantity is defined from the finite subset of nonpositive powers of the Laurent series for u .

Definition 20. Given a family (p, u_0) , the *singular part operator* \mathcal{D} is defined as

$$\text{Log } \varphi \mapsto \mathcal{D} \text{Log } \varphi = u_T(0) - u_T(\infty), \quad (83)$$

in which the notation $u_T(\varphi_0)$, which emphasizes the dependence on the movable constant φ_0 , stands for the principal part (T like truncation) of the Laurent series (55), i.e. the finite subset of its nonpositive powers

$$u_T(\varphi_0) = \sum_{j=0}^{-p} u_j \chi^{j+p}. \tag{84}$$

For most PDEs, this operator is linear. For the Laurent series already considered (63), (72), (73), (80), the operator is, respectively, $\mathcal{D} = -2a\partial_x^2, a, a\partial_x, d\partial_x$. For the Kuramoto-Sivashinsky equation (3), there exists a unique Laurent series (55) with $p = -3$ (given by (5) for a particular value of χ , and by the derivative of (347) for any χ), with a singular part operator equal to

$$\mathcal{D} = 60\nu\partial_x^3 + (60/19)\mu\partial_x. \tag{85}$$

This is precisely the third order linear operator on the rhs of (49).

4.3 The various ways to pass or fail the Painlevé test for PDEs

If one processes a multidimensional PDE the coefficients of which depend on some parameters μ ,

$$\mathbf{E}(\mathbf{u}, \mathbf{x}; \mu) = 0, \tag{86}$$

(boldface means multicomponent), the Painlevé test generates the following output :

1. leading order (p, u_0) , Fuchs indices i and singular part operator \mathcal{D} for each admissible family,
2. diophantine conditions that all singularity orders p and all Fuchs indices i be integral, conditions whose solution creates constraints of the type

$$F(\mu, C) = 0, \tag{87}$$

3. no-log conditions

$$\forall i \forall n \forall u_j : \mathbf{Q}_i^n(\mu, S, C, u_j) = 0, \tag{88}$$

arising from any integral Fuchs index i , in which n is the Fuchsian perturbation order [28] if necessary, u_j are the arbitrary coefficients u_{arb} introduced at earlier Fuchs indices j .

In particular, the Laurent series (55) are of no use and should not be computed beyond the highest Fuchs integer. All this output (items 1 and 3) is easily produced with a computer algebra program and, in all further examples, we will simply list these results without any more detail.

Strictly speaking, the answer provided by the test to the question “Has the PDE the PP?” is either *no* (at least one of the necessary conditions fails) or *maybe* (all necessary conditions are satisfied, and the PDE may possess the PP but this still has to be proven). It is never *yes*, as shown by the famous counterexample of Painlevé (the second order ODE with the general solution $\wp(\lambda \text{Log}(c_1x + c_2), g_2, g_3)$, which therefore has the PP iff $2\pi i\lambda$ is a period of the Weierstrass elliptic function \wp , a transcendental condition impossible to generate by a finite algebraic procedure).

Now that the necessary part (i.e. the Painlevé test) of Painlevé analysis is finished, let us turn to the question of sufficiency.

To reach our goal which is to obtain as many analytic results as possible, we do not adopt such a drastic point of view, but the opposite one. Instead of the logical *and* performed by the mathematician on all the necessary conditions generated by the test, we perform a logical *or* operation on these conditions. Therefore the above Painlevé test must be performed to its end, i.e. without stopping even in case of failure of some condition, so as to collect all the necessary conditions. Turning to sufficiency, these conditions have to be examined independently in the hope of finding some global element of integrability. An application of this point of view to the Lorenz model, a third order ODE, can be found in Section 6.7 of Ref. [26].

If the PDE under study possesses a singlevalued exact solution, there must exist a Laurent series (55) which represents it locally. Therefore the practical criterium to be implemented deals with the existence of *particular* Laurent series, and the result of the test belongs to one of the following mutually exclusive situations.

1. (The best situation) Success of the test, at least for some values of μ selected by the test. The PDE may have the PP, and one must look for its BT;
2. There exists at least one value of $(\mu, \varphi, u_{\text{arb}})$ which ensures the existence of a particular Laurent series. For these values, an exact solution may exist;
3. There exists at least one value of $(\mu, \varphi, u_{\text{arb}})$ enforcing some of, but not all, the no-log conditions of at least one particular Laurent series. Quite probably no exact solution exists, but there may exist a conservation law (a first integral for an ODE);
4. (The worst situation) There is no value of $(\mu, \varphi, u_{\text{arb}})$ enforcing at least one of the no-log conditions of the various series. Quite probably the PDE is chaotic and possesses no exact solution at all.

Examples of these various situations are, respectively :

1. All the PDEs which have the PP (sine-Gordon, Korteweg-de Vries, ...), but also the counterexample of Painlevé quoted above;
2. The equation of Kuramoto and Sivashinsky (3), with the particular Laurent series (5);

3. The Lorenz model for $b = 2\sigma$, for which the no-log condition at $i = 4$ is violated and there exists a first integral;
4. The Rössler dynamical system for which the unique family has the never satisfied condition $Q_2 \equiv 16 = 0$.

5 Ingredients of the “singular manifold method”

The methods to handle the integrable and nonintegrable situations are the same, simply a more or less important result is obtained.

The goal is to find a (possibly degenerate) couple (Darboux transformation, Lax pair) in order to deduce the Bäcklund transformation or, if a BT does not exist, to generate some exact solutions.

The full Laurent series is of no help, for this is not an exact solution according to the definition in Section 2. Since this is the only available piece of information and since a finite (closed form) expression is required to represent an exact solution, let us represent, and this is the idea of Weiss, Tabor and Carnevale [119], an unknown exact solution u as the sum of a singular part, built from the finite principal part of the Laurent series (i.e. the finite number of terms with negative powers), and of a regular part made of one term denoted U . This assumption is identical to that of a Darboux transformation (46), in which nothing would be specified on U .

This method is widely known as the *singular manifold method* or *truncation method* because it selects the beginning of the Laurent series and discards (“truncates”) the remaining infinite part.

Since its introduction by WTC [119], it has been improved in many directions [86,43,53,88,34,101,90], and we present below the current status of the method.

5.1 The ODE situation

For the six ordinary differential equations (ODE) (P1)–(P6) which bear his name, Painlevé proved the PP by showing [97,98] the existence of one (case of (P1)) or two ((P2)–(P6)) function(s) $\tau = \tau_1, \tau_2$, called tau-functions, linked to the general solution u by logarithmic derivatives

$$(P1) : u = \mathcal{D}_1 \text{Log } \tau \tag{89}$$

$$(Pn), n = 2, \dots, 6 : u = \mathcal{D}_n(\text{Log } \tau_1 - \text{Log } \tau_2) \tag{90}$$

where the operators \mathcal{D}_n are linear:

$$\mathcal{D}_1 = -\partial_x^2, \mathcal{D}_2 = \mathcal{D}_4 = \pm \partial_x, \mathcal{D}_3 = \pm e^{-x} \partial_x, \tag{91}$$

$$\mathcal{D}_5 = \pm x e^{-x} (2\alpha)^{-1/2} \partial_x, \mathcal{D}_6 = \pm x(x-1) e^{-x} (2\alpha)^{-1/2} \partial_x. \tag{92}$$

These functions τ_1, τ_2 satisfy third order nonlinear ODEs and they have the same kind of singularities than solutions of *linear* ODEs, namely they have

no movable singularities at all; they are entire functions for (P1)–(P5), and their only singularities for (P6) are the three fixed critical points $(\infty, 0, 1)$.

ODEs cannot possess an auto-BT, since the number of independent arbitrary coefficients in a solution cannot exceed the order of the ODE. They can however possess a Schlesinger transformation (see definition Section 11).

5.2 Transposition of the ODE situation to PDEs

For PDEs, similar ideas prevail. The analogue of (89)–(90), with an additional rhs U , is now the Darboux transformation (46), and the scalar(s) ψ to which the scalar(s) τ are linked by (47) are assumed to satisfy a linear system, the Lax pair.

Another interesting observation must be made. There seems to exist two and only two classes of integrable $1 + 1$ -dimensional PDEs, at least at the level of the base member of a hierarchy : those which have only one family of movable singularities, and those which have only pairs of families with opposite principal parts, similarly to the distinction between (P1) on one side and (P2)–(P6) on the other side. Among the $1 + 1$ -dimensional integrable equations, those with one family include KdV, the AKNS, Hirota-Satsuma and Boussinesq equations; they also include the Sawada-Kotera, Kaup-Kupershmidt and Tzitzéica equations because only one of their two families is relevant [90,37]. Equations with pairs of opposite families include sine-Gordon, mKdV and Broer-Kaup (two families each), NLS (four families).

5.3 The singular manifold method as a Darboux transformation

As qualitatively described in Section 5, the singular manifold method looks very much like a resummation of the Laurent series, just like the geometric series

$$\sum_{j=0}^{+\infty} x^j, \quad x \rightarrow 0, \quad (93)$$

becomes a finite sum in the resummation variable $X = x/(1 - x)$

$$\sum_{J=0}^1 X^J, \quad X \rightarrow 0. \quad (94)$$

The principle of the method is the following [119]. One first notices that the (infinite) Laurent series (55) in the variable $\varphi - \varphi_0$ can be rewritten as the sum of two terms

$$u = \mathcal{D} \text{Log } \tau + \text{regular part}. \quad (95)$$

The first term $\mathcal{D} \operatorname{Log} \tau$, built from the singular part operator defined in Section 4.2, is a finite Laurent series and, if τ is any variable fulfilling the two requirements for an expansion variable enunciated in Section 4.1, it captures all the singularities of u when $\varphi \rightarrow \varphi_0$. The second term, temporarily called “regular part” for this reason, is yet unspecified. The sum of these two terms is therefore a finite Laurent series (hence the name *truncated series*), and the variable τ is a *resummation variable* which has made the former infinite series in $\varphi - \varphi_0$ a finite one. One then tries to identify this resummation (95) with the definition of a Darboux transformation (46). This involves two features. The first feature is to uncover a link (47) between τ and a scalar component ψ of a Lax pair. The second feature is to prove that the left over “regular part” is indeed a second solution to the PDE under study.

5.4 The degenerate case of linearizable equations

The Burgers equation (71), under the transformation of Forsyth (Ref. [48] p. 106),

$$v = a \operatorname{Log} \tau, \quad \tau = \psi, \quad (96)$$

is linearized into the heat equation

$$b\psi_t + \psi_{xx} + G(t)\psi = 0. \quad (97)$$

This can be considered as a degenerate Darboux transformation (46), in which U is identically zero and ψ satisfies a single linear equation, not a pair of linear equations, so this fits the general scheme.

Another classical example is the second order particular Monge-Ampère equation $s + pq = 0$, linearized into the d’Alembert equation $s = 0$:

$$s + pq \equiv u_{xt} + u_x u_t = 0, \quad (98)$$

$$u = \operatorname{Log} \tau, \quad \tau = \psi, \quad \psi_{xt} = 0. \quad (99)$$

5.5 Choices of Lax pairs and equivalent Riccati pseudopotentials

To fix the ideas, we list here a few usual second order and third order Lax pairs depending on undetermined coefficients, together with the constraints imposed on these coefficients by the commutativity condition.

It is sometimes appropriate to represent an n -th order Lax pair by the $2(n - 1)$ equations satisfied by an equivalent $(n - 1)$ -component pseudopotential \mathbf{Y} of Riccati type, the first component of which is chosen as

$$Y_1 = \psi_x / \psi, \quad (100)$$

in which ψ is a scalar component of the Lax pair.

Second-order Lax pairs and their privilege

The general second-order scalar Lax pair reads, in the case of two independent variables (x, t) ,

$$L_1\psi \equiv \psi_{xx} - d\psi_x - a\psi = 0, \quad (101)$$

$$L_2\psi \equiv \psi_t - b\psi_x - c\psi = 0, \quad (102)$$

$$[L_1, L_2] \equiv X_0 + X_1\partial_x, \quad (103)$$

$$(104)$$

$$X_0 \equiv -a_t + a_x b + 2ab_x + c_{xx} - c_x d = 0, \quad (105)$$

$$X_1 \equiv -d_t + (b_x + 2c - bd)_x = 0. \quad (106)$$

For the inverse scattering method to apply, the coefficients (d, a) of the x -part (101) are required to depend linearly on the field U of the PDE.

The Lax pair (101)–(102) is identical to a linearized version of the Riccati system satisfied by the most general expansion variable Y defined by (66), under the correspondence

$$Y = B^{-1} \frac{\psi}{\psi_x}, \quad B \neq 0, \quad (107)$$

$$d = 2A, \quad a = A_x - A^2 - S/2, \quad b = -C, \quad c = C_x/2 + AC + \int A_t dx, \quad (108)$$

and the commutator of the Lax pair is (62).

In particular, when the coefficient d is zero or when, by a linear change $\psi \mapsto e^{\int d dx/2} \psi$, it can be set to zero without altering the linearity of a on U , the correspondence is [86]

$$\chi = \frac{\psi}{\psi_x}, \quad B = 1, \quad A = 0, \quad (109)$$

$$d = 0, \quad a = -S/2, \quad b = -C, \quad c = C_x/2, \quad (110)$$

$$L_1\psi \equiv \psi_{xx} + \frac{S}{2}\psi = 0, \quad (111)$$

$$L_2\psi \equiv \psi_t + C\psi_x - \frac{C_x}{2}\psi = 0, \quad (112)$$

$$2[L_1, L_2] \equiv X = S_t + C_{xxx} + CS_x + 2C_x S = 0. \quad (113)$$

Therefore second order Lax pairs are privileged in the singularity approach, in the sense that their coefficients can be identified with the elementary homographic invariants S, C of the invariant Painlevé analysis and, if appropriate, A, B . Conversely, and this has historically been the reason of some errors described in Section 8.2, at the stage of searching for the BT, these homographic invariants S, C are useless when the Lax order is higher than two and they should not be considered.

As explained in Section 5.3, given a Lax pair, one should define from it either one or two scalars ψ_f . Consider the second order Lax pair defined by the gradient of Y . Then, for one-family PDEs, this unique scalar ψ is defined by (107). For two-family PDEs, the two scalars ψ_f are defined by

$$Y = \frac{\psi_1}{\psi_2}, \quad (114)$$

which leads to the zero-curvature representation of the Lax pair

$$(\partial_x - L) \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = 0, \quad L = \begin{pmatrix} -A - B^{-1}B_x/2 & B^{-1} \\ B(A_x - A^2 - S/2) & A + B^{-1}B_x/2 \end{pmatrix}, \quad (115)$$

$$(\partial_t - M) \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = 0, \quad (116)$$

$$M = \begin{pmatrix} AC + C_x/2 - B^{-1}B_t/2 & -CB^{-1} \\ B((CS + C_{xx})/2 + A_t + CA^2 + C_xA) & -AC - C_x/2 + B^{-1}B_t/2 \end{pmatrix}.$$

The reason why the Riccati form is the most suitable characterization of the Lax pair is that it allows two linearizations [88,101], namely (107) and (114), depending on whether the PDE has one family or two opposite families.

Third-order Lax pairs

The general third-order scalar Lax pair is defined as

$$L_1\psi \equiv \psi_{xxx} - f\psi_{xx} - a\psi_x - b\psi = 0, \quad (117)$$

$$L_2\psi \equiv \psi_t - c\psi_{xx} - d\psi_x - e\psi = 0, \quad (118)$$

$$[L_1, L_2] \equiv X_0 + X_1\partial_x + X_2\partial_x^2, \quad (119)$$

$$\begin{aligned} X_0 \equiv & -b_t - ae_x + e_{xxx} + b_{xx}c + 2bcf_x + bc_xf - e_{xx}f \\ & + 3bc_{xx} + 3b_xc_x + 3bd_x + b_xd = 0, \end{aligned} \quad (120)$$

$$\begin{aligned} X_1 \equiv & -a_t + 3e_{xx} + 2b_xc + a_{xx}c + d_{xxx} + 3ac_{xx} + 2ad_x \\ & + 3a_xc_x + 3bc_x + a_xd + 2acf_x + ac_xf - 2e_xf - d_{xx}f = 0, \end{aligned} \quad (121)$$

$$X_2 \equiv -f_t + (cf^2 + cf_x + 2c_xf + df + 2ac + c_{xx} + 3d_x + 3e)_x = 0. \quad (122)$$

An equivalent two-component pseudopotential is the projective Riccati one $\mathbf{Y} = (Y_1, Y_2)$ [86,87] (written below, for simplicity, in the case $f = 0$)

$$Y_1 = \frac{\psi_x}{\psi}, \quad Y_2 = \frac{\psi_{xx}}{\psi}, \quad (123)$$

$$Y_{1,x} = -Y_1^2 + Y_2, \quad (124)$$

$$Y_{2,x} = -Y_1Y_2 + aY_1 + b, \quad (125)$$

$$Y_{1,t} = -(dY_1 + cY_2)Y_1 + (ac + d_x)Y_1 + (c_x + d)Y_2 + e_x + bc \quad (126)$$

$$= (cY_2 + dY_1 + e)_x, \quad (127)$$

$$Y_{2,t} = -(dY_1 + cY_2)Y_2 + (2ac_x + a_xc + bc + d_{xx} + ad + 2e_x)Y_1$$

$$+(c_{xx} + 2d_x + ac)Y_2 + 2bc_x + b_xc + bd + e_{xx}, \quad (128)$$

$$Y_{1,tx} - Y_{1,xt} = X_1 + X_2Y_1, \quad (129)$$

$$Y_{2,tx} - Y_{2,xt} = -X_0 + X_2Y_2. \quad (130)$$

When there is no reason to distinguish between x and t , for instance because the PDE is invariant under the permutation (Lorentz transformation)

$$\mathcal{P} : (\partial_x, \partial_t) \rightarrow (\partial_t, \partial_x), \quad (131)$$

it is natural to consider the following third-order matrix Lax pair, invariant under (131), defined in the basis (ψ_x, ψ_t, ψ) [37],

$$(\partial_x - L) \begin{pmatrix} \psi_x \\ \psi_t \\ \psi \end{pmatrix} = 0, \quad L = \begin{pmatrix} f_1 & f_2 & f_3 \\ g_1 & g_2 & g_3 \\ 1 & 0 & 0 \end{pmatrix}, \quad (132)$$

$$(\partial_t - M) \begin{pmatrix} \psi_x \\ \psi_t \\ \psi \end{pmatrix} = 0, \quad M = \begin{pmatrix} g_1 & g_2 & g_3 \\ h_1 & h_2 & h_3 \\ 0 & 1 & 0 \end{pmatrix}. \quad (133)$$

In the equivalent projective Riccati components (Y_1, Y_2)

$$Y_1 = \frac{\psi_x}{\psi}, \quad Y_2 = \frac{\psi_t}{\psi}, \quad (134)$$

with the property $Y_{1,t} = Y_{2,x}$, it is defined as

$$Y_{1,x} = -Y_1^2 + f_1Y_1 + f_2Y_2 + f_3, \quad (135)$$

$$Y_{2,x} = -Y_1Y_2 + g_1Y_1 + g_2Y_2 + g_3, \quad (136)$$

$$Y_{1,t} = -Y_1Y_2 + g_1Y_1 + g_2Y_2 + g_3, \quad (137)$$

$$Y_{2,t} = -Y_2^2 + h_1Y_1 + h_2Y_2 + h_3. \quad (138)$$

The nine functions $f_j, g_j, h_j, j = 1, 2, 3$, must satisfy six cross-derivative conditions $X_j = 0$

$$(Y_{1,x})_t - (Y_{1,t})_x = X_0 + X_1Y_1 + X_2Y_2 = 0, \quad (139)$$

$$(Y_{2,x})_t - (Y_{2,t})_x = X_3 + X_4Y_1 + X_5Y_2 = 0, \quad (140)$$

easy to write explicitly. It is worth noticing that there exists no such invariant second-order matrix Lax pair.

5.6 The admissible relations between τ and ψ

By elimination of ∂_t , one of the two PDEs defining the BT to be found can be made an ODE, e.g. (64) or (152). This nonlinear ODE, with coefficients depending on U and, in the 1+1-dimensional case, on an arbitrary constant λ , has the property [90] of being linearizable. This very strong property restricts the admissible choices (47) to a finite number of possibilities, and full details can be found in Musette lecture [85].

6 The algorithm of the singular manifold method

We now have all the ingredients to give a general exposition of the method in the form of an algorithm. The present exposition largely follows the lines of Ref. [90].

The various situations thus implemented are : one-family and two-opposite-family PDEs, second or higher order Lax pair, various allowed links between the two sets of functions (τ, ψ) .

Consider a PDE (53) with only one family of movable singularities or exactly two families of movable singularities with opposite values of u_0 , and denote \mathcal{D} the *singular part operator* of either the unique family or any one of the two opposite families.

First step. Assume a Darboux transformation defined as

$$u = U + \mathcal{D}(\text{Log } \tau_1 - \text{Log } \tau_2), \quad E(u) = 0, \quad (141)$$

with u a solution of the PDE under consideration, U an unspecified field which most of the time will be found to be a second solution of the PDE, τ_f the “entire” function (or more precisely ratio of entire functions) attached to each family f . For one-family PDEs, one denotes $\tau_1 = \tau, \tau_2 = 1$, so the DT assumption (141) becomes

$$u = U + \mathcal{D} \text{Log } \tau, \quad E(u) = 0. \quad (142)$$

A consequence of the assumption (141) is the existence of the involution

$$\forall f : (u, U, \tau_f) \mapsto (U, u, \tau_f^{-1}), \quad (143)$$

since the operator \mathcal{D} is linear, and, for two-family PDEs, of the involution

$$\forall (u, U) : (\mathcal{D}, \tau_1, \tau_2) \mapsto (-\mathcal{D}, \tau_2, \tau_1). \quad (144)$$

Second step. Choose the order two, then three, then \dots , for the unknown Lax pair, and define one or two (as many as the number of families) *scalars* ψ_f from the component(s) of its wave vector (e.g. the scalar wave vector if the PDE has one family and the pair is defined in scalar form). Such sample Lax pairs and scalars can be found in Section 5.5.

Third step. Choose an explicit link F

$$\forall f : \mathcal{D} \text{Log } \tau_f = F(\psi_f), \quad (145)$$

the same for each family f , between the functions τ_f and the scalars ψ_f defined from the Lax pair. According to Section 5.6, at each scattering order, there exists only a finite number of choices (145), among them the most frequent one

$$\forall f : \tau_f = \psi_f. \quad (146)$$

Fourth step. Define the “truncation” and solve it, that is to say : with the assumptions (141) for a DT, (145) for a link between τ_f and ψ_f , (101)–(102) or (117)–(118) or other for the Lax pair in ψ , express $E(u)$ as a polynomial in the derivatives of ψ_f which is irreducible *modulo* the Lax pair. For the above pairs and a one-family PDE, this amounts to eliminate any derivative of ψ of order in (x, t) higher than or equal to $(2, 0)$ or $(0, 1)$ (second order case) or to $(3, 0)$ or $(0, 1)$ (third order), thus resulting in a polynomial of one variable ψ_x/ψ (second order) or two variables $\psi_x/\psi, \psi_{xx}/\psi$ (third order)

$$E(u) = \sum_{j=0}^{-q} E_j(S, C, U)(\psi/\psi_x)^{j+q} \text{ (one-family PDE, second order), (147)}$$

$$E(u) = \sum_{k \geq 0} \sum_{l \geq 0} E_{k,l}(a, b, c, d, e, U)(\psi_x/\psi)^k (\psi_{xx}/\psi)^l$$

(one-family PDE, third order). (148)

Since one has no more information on this polynomial $E(u)$ except the fact that it must vanish, one requests that it identically vanishes, by solving the set of *determining equations*

$$\forall j \quad E_j(S, C, U) = 0 \text{ (one-family PDE, second order) (149)}$$

$$\forall k \forall l \quad E_{k,l}(a, b, c, d, e, U) = 0 \text{ (one-family PDE, third order) (150)}$$

for the unknown coefficients (S, C) or (a, b, c, d, e) as functions of U , and one establishes the constraint(s) on U by eliminating (S, C) or (a, b, c, d, e) . The strategy of resolution is developed in Section 7.3.

The constraints on U reflect the integrability level of the PDE. If the only constraint on U is to satisfy some PDE, one is on the way to an auto-BT if the PDE for U is the same as the PDE for u , or to a remarkable correspondence (hetero-BT) between the two PDEs.

The second, third and fourth steps must be repeated until a success occurs. The process is successful if and only if all the following conditions are met

1. U comes out with one constraint exactly, namely : to be a solution of some PDE,
2. (if an auto-BT is desired) the PDE satisfied by U is identical to (53),
3. the vanishing of the commutator $[L_1, L_2]$ is equivalent to the vanishing of the PDE satisfied by U ,
4. in the 1+1-dimensional case only and if the PDE satisfied by U is identical to (53), the coefficients depend on an arbitrary constant λ , the spectral or Bäcklund parameter.

At this stage, one has obtained the DT and the Lax pair.

Fifth step. Obtain the two equations for the BT by eliminating ψ_f [14] between the DT and the Lax pair. This sometimes uneasy operation when the order n of the Lax pair is too high may become elementary by considering

the equivalent Riccati representation of the Lax pair and eliminating the appropriate components of \mathbf{Y} rather than ψ . Assume for instance that $\tau = \psi$, $\mathcal{D} = \partial_x$, and the PDE has only one family. Then Eq. (141) reads

$$Y_1 = u - U \quad (151)$$

with Y_1 defined in (100), and the BT is computed as follows : eliminate all the components of \mathbf{Y} but Y_1 between the equations for the gradient of \mathbf{Y} , then in the resulting equations substitute Y_1 as defined in (151).

If the computation of the BT requires the elimination of Y_2 between (124)–(128), this BT is

$$Y_{1,xx} + 3Y_1Y_{1,x} + Y_1^3 - aY_1 - b = 0, \quad (152)$$

$$Y_{1,t} - (cY_{1,x} + cY_1^2 + dY_1 + e)_x = 0, \quad (153)$$

$$(Y_{1,xx})_t - (Y_{1,t})_{xx} = X_0 + X_1Y_1 + X_2Y_1^2 = 0, \quad (154)$$

in which Y_1 is replaced by an expression of $u - U$, e.g. (151).

Although, let us repeat it, the method equally applies to integrable as well as nonintegrable PDEs, examples are split according to that distinction, to help the reader to choose his/her field of interest.

6.1 Where to truncate, and with which variable?

This Section is self-contained, and mainly destined to persons accustomed to perform the WTC truncation. Although some paragraphs might be redundant with Section 6, it may help the reader by presenting a complementary point of view.

Let us assume in this Section that the unknown Lax pair is second order. Then the truncation defined in the fourth step of Section 6 is performed in the style of Weiss *et al.* [119], i.e. with a single variable. This WTC truncation consists in forcing the series (55) to terminate; let us denote p and q the singularity orders of u and $E(u)$, $-p'$ the rank at which the series for u stops, and $-q'$ the corresponding rank of the series for E

$$u = \sum_{j=0}^{-p'} u_j Z^{j+p}, \quad u_0 u_{-p'} \neq 0, \quad E = \sum_{j=0}^{-q'} E_j Z^{j+q}, \quad (155)$$

in which the *truncation variable* Z chosen by WTC is $Z = \varphi - \varphi_0$. Since one has no more information on Z , the method of WTC is to require the separate satisfaction of each of the *truncation equations*

$$\forall j = 0, \dots, -q' : E_j = 0. \quad (156)$$

In earlier presentations of the method, one had to prove by recurrence that, assuming that enough consecutive coefficients u_j vanish beyond $j = -p'$,

then all further coefficients u_j would vanish. This painful task is useless if one defines the process as done above.

The first question to be solved is : what are the admissible values of p' , i.e. those which respect the condition $u_{-p'} \neq 0$?

The answer depends on the choice of the truncation variable Z . In Section 4.1, three choices were presented, $Z = \varphi - \varphi_0$, χ or Y , respectively defined by equations (54), (58), (66), with the property that any two of their inverse are linearly dependent.

The advantage of χ or Y over $\varphi - \varphi_0$ is the following. The gradient of χ (resp. Y) is a polynomial of degree two in χ (resp. Y), so each derivation of a monomial aZ^k increases the degree by one, while the gradient of $\varphi - \varphi_0$ is a polynomial of degree zero in $\varphi - \varphi_0$, so each derivation decreases the degree by one. Consequently, one finds two solutions and only two to the condition $u_{-p'} \neq 0$ [100]

1. $p' = p, q' = q$, in which case the three truncations are identical, since the three sets of equations $E_j = 0$ are equivalent (the finite sum $\sum E_j Z^{j+q}$ is just the same polynomial of Z^{-1} written with three choices for its base variable),
2. for χ and Y only, $p' = 2p, q' = 2q$, in which case the two truncations are different since the two sets of equations $E_j = 0$ are inequivalent (they are equivalent only if $A = 0$).

To perform the first truncation $p' = p, q' = q$, one must then choose $Z = \chi$ since Y brings no more information and $\varphi - \varphi_0$ creates equivalent but lengthier expressions.

To perform the second truncation $p' = 2p, q' = 2q$, one must choose $Z = Y$, since χ would create the *a priori* constraint $A = 0$.

The second question to be solved is : given some PDE with such and such structure of singularities, and assuming that one of the above two truncations is relevant (which is a separate topic), which one should be selected?

The answer lies in the two elementary identities [31]

$$\tanh z - \frac{1}{\tanh z} = -2i \operatorname{sech} \left[2z + i\frac{\pi}{2} \right], \quad \tanh z + \frac{1}{\tanh z} = 2 \tanh \left[2z + i\frac{\pi}{2} \right]. \quad (157)$$

Let us explain why on two examples, the ODEs whose general solution is $\tanh(x - x_0)$ and $\operatorname{sech}(x - x_0)$, namely

$$E \equiv u' + u^2 - 1 = 0, \quad u = \tanh(x - x_0), \quad (158)$$

$$E \equiv v'^2 + a^{-2}v^4 - v^2 = 0, \quad v = a \operatorname{sech}(x - x_0), \quad (159)$$

(this is just for convenience that we do not set $a = 1$). Equation (158) has the single family

$$p = -1, q = -2, u_0 = 1, \text{ Fuchs indices} = (-1), \quad (160)$$

and equation (159) has the two opposite families

$$p = -1, q = -4, v_0 = ia, \text{ Fuchs indices} = (-1), \quad (161)$$

in which ia denotes any square root of $-a^2$. The first truncation

$$u = \sum_{j=0}^{-p} u_j \chi^{j+p}, \quad E = \sum_{j=0}^{-q} E_j \chi^{j+q}, \quad \forall j : E_j = 0, \quad (162)$$

generates the respective results

$$u = \chi^{-1}, \quad S = -2, \quad (163)$$

$$v = ia\chi^{-1}, \quad E_2 \equiv a^2(1 - S) = 0, \quad E_3 \equiv 0, \quad E_4 \equiv -a^2 S^2/4, \quad (164)$$

thus providing (after integration of the Riccati ODE (64)) the general solution of equation (158), and no solution at all for equation (159).

The second truncation

$$u = \sum_{j=0}^{-2p} u_j Y^{j+p}, \quad E = \sum_{j=0}^{-2q} E_j Y^{j+q}, \quad \forall j : E_j = 0, \quad (165)$$

generates the respective results

$$u = B^{-1}Y^{-1} + (1/4)BY, \quad A = 0, \quad S = -1/2, \quad B \text{ arbitrary}, \quad (166)$$

$$v = iaB^{-1}Y^{-1} - (1/4)iaBY, \quad A = 0, \quad S = -1/2, \quad B \text{ arbitrary}, \quad (167)$$

thus providing, thanks to the identities (157), the general solution for both equations.

The conclusions from this exercise which can be generalized are :

1. for PDEs with only one family, the second truncation brings no additional information as compared to the first one and is always useless;
2. for PDEs with two opposite families (two opposite values of u_0 for a same value of p), the first truncation can never provide the general solution and can only provide particular solutions, while the second one may provide the general solution.

This defines the guideline to be followed in the respective Sections 7 and 9. The question of the relevance of the parameter B , which seems useless in the above two examples, is addressed in Section 9.

7 The singular manifold method applied to one-family PDEs

7.1 Integrable equations with a second order Lax pair

There is only one truncation variable, which must be chosen as χ .

Weiss introduced a nice notion, initially for one-family integrable equations with a second order Lax pair, later extended to two-family such equations by Pickering [101]. This is the following.

Definition 21. ([20]) Consider the set of $-q+p$ *determining equations* (149) $E_j = 0$, which depend on (S, C, U) . One calls **singular manifold equation** (SME) the result of the elimination of U between them.

In the two-family situation, these determining equations also depend on (A, B) , see (165), and the extension of this definition [101] is to also require the elimination of (A, B) .

Despite its name, originally restricted to integrable equations, the SME can be made of several equations in the nonintegrable case.

The SME has the following properties.

1. unicity, whatever be the integrability of the PDE,
2. invariance under homography by construction [19], i.e. dependence only on one Schwarzian S and as many C quantities as independent variables other than the one in the Schwarzian,
3. the SME set is made of one and only one equation if and only if the PDE is integrable.

Although one can define a SME whatever be the order of the Lax pair, it is inconsistent, as will be explained in Section 8.2, to do so whenever this order is higher than two.

The Liouville equation

It is convenient to consider, following Zhiber and Shabat [125], the equation

$$E(u) \equiv u_{xt} + \alpha e^u + a_1 e^{-u} + a_2 e^{-2u} = 0, \quad \alpha \neq 0, \quad (168)$$

which has the advantage to include the Liouville equation $a_1 = a_2 = 0$, the sine-Gordon equation ($a_1 \neq 0, a_2 = 0$) and the Tzitzéica equation ($a_1 = 0, a_2 \neq 0$). As to the case $a_1 a_2 \neq 0$, it fails the test. Let us consider here the Liouville case. The results to be found are its auto-BT [81] and its hetero-BT to the d'Alembert equation. This will be achieved with two different truncations.

Although not algebraic in u , the PDE is algebraic in either e^u or e^{-u} .

Equation (168) always possesses the family

$$e^u \sim -(2/\alpha)\varphi_x\varphi_t(\varphi - \varphi_0)^{-2}, \quad \text{indices } (-1, 2), \quad \mathcal{D} = (2/\alpha)\partial_x\partial_t. \quad (169)$$

For Liouville, this is the only family.

The special form of Liouville equation allows the assumption

$$e^u = \mathcal{D} \operatorname{Log} \tau + e^U, \quad E(u) = 0, \quad \mathcal{D} = (2/\alpha) \partial_x \partial_t, \quad (170)$$

to be integrated twice to yield

$$u = -2 \operatorname{Log} \tau + V, \quad E(u) = \sum_{j=0}^2 E_j \tau^{j-2} = 0, \quad (171)$$

in which nothing is assumed on V .

The Liouville equation is nongeneric for the singular manifold method in the sense that it is linearizable into another equation (thus, it should even *not* be part of the Section 7.1).

Therefore we define the first truncation in an exceptional way, namely we do not assume any linear relations on $\tau \equiv \psi$ and just treat τ as the truncation variable. The three determining equations are then quite simple [37]

$$E_0 \equiv 2\tau_x \tau_t + \alpha e^V = 0, \quad (172)$$

$$E_1 \equiv \tau_{xt} = 0, \quad (173)$$

$$E_2 \equiv V_{xt} = 0, \quad (174)$$

and their general solution depends on two arbitrary functions of one variable

$$\tau = f(x) + g(t), \quad (175)$$

$$e^V = -\frac{2}{\alpha} \tau_x \tau_t = -\frac{2}{\alpha} f'(x) g'(t), \quad (176)$$

$$e^u = -\frac{2}{\alpha} \frac{\tau_x \tau_t}{\tau^2} = -\frac{2}{\alpha} \frac{f'(x) g'(t)}{(f(x) + g(t))^2}, \quad (177)$$

$$e^U = \tau^{-2} e^V + \frac{2}{\alpha} \frac{\tau_x \tau_t}{\tau^2} = 0. \quad (178)$$

Thus, the two fields u and V are the general solution of, respectively, the Liouville and d'Alembert equations. The hetero-BT between these two equations is provided by the elimination of f and g between (176), (177) and the x - and t -derivatives of (171)

$$(u - v)_x = \alpha \lambda e^{(u+v)/2}, \quad (179)$$

$$(u + v)_t = -2\lambda^{-1} e^{(u-v)/2}, \quad (180)$$

in which v is another solution of d'Alembert equation defined by

$$e^v = (\lambda \tau_t)^{-2} e^V = -\frac{2}{\alpha} \lambda^{-2} \frac{f'(x)}{g'(t)}. \quad (181)$$

Remark. When performing the truncation (170), Tamizhmani and Lakshmanan [109] already found $e^U = 0, \tau_{xt} = 0$ as a *particular* solution, while

the above truncation (171) proves it to be the *general* solution. Another difference between the two truncations is the presence of a field V in (171), which allows us to find in addition the hetero-BT between the Liouville and d'Alembert equations.

Let us now define the second truncation, by the assumption

$$u = -2 \operatorname{Log} \tau + \tilde{W}, \quad (182)$$

and the link (146), with ψ solution of the Lax pair (101)–(102). Introducing the Riccati variable Y defined by (107), this second truncation is equivalent to [37]

$$\begin{aligned} u &= -2 \operatorname{Log} Y + W, \quad Y^{-1} = B(\chi^{-1} + A), \\ E(u) &= \sum_{j=0}^4 E_j(S, C, A, B, W) Y^{j-2}, \quad \forall j: E_j = 0, \end{aligned} \quad (183)$$

and its result is recovered from the truncation of sine-Gordon in Section 9.1 by simply setting $a_1 = 0$.

The AKNS equation

The AKNS equation [2]

$$E(u) \equiv u_{xxxt} + 4\alpha^{-1}(2(u_x - \beta)u_{xt} + (u_t - \gamma)u_{xx}) = 0 \quad (184)$$

admits the single family

$$p = -1, \quad q = -5, \quad u_0 = \alpha, \quad \text{indices } (-1, 1, 4, 6), \quad \mathcal{D} = \alpha \partial_x, \quad (185)$$

so the assumption for the DT is (142). Let us choose at second step the scalar Lax pair (111)–(112) for ψ , at third step the link (146) between τ and ψ . Then there are only three non identically zero determining equations (149) [83]

$$E_2 \equiv 4\alpha SC + 8(U_t - \gamma) - 16C(U_x - \beta) = 0, \quad (186)$$

$$E_3 \equiv -\alpha(CS_x + 4SC_x) + 16C_x(U_x - \beta) - 8U_{xt} + 4CU_{xx} = 0, \quad (187)$$

$$\begin{aligned} E_5 \equiv & E(U) + (\alpha/2)(2SS_t - CSS_x - S_{xxt} - S_x C_{xx}) - 2S_x(U_t - \gamma) \\ & - 4S_t(U_x - \beta) - 4SU_{xt} + 2(SC + C_{xx})U_{xx} = 0, \end{aligned} \quad (188)$$

plus the ever present condition $X = 0$, Eqn. (113). Their detailed resolution for (U_x, U_t) is as follows. One eliminates U_t between E_2 and E_3

$$E_3 + E_{2,x} \equiv 3C(-4U_{xx} + \alpha S_x) = 0, \quad (189)$$

discards the nongeneric solution $C = 0, U_t = \gamma, S_t = 0$, introduces an arbitrary function of t after one integration, and solves for U_x

$$U_x - \beta = (\alpha/4)(S + 2\lambda(t)). \quad (190)$$

Then E_2 is solved for U_t

$$U_t - \gamma = \alpha\lambda(t)C. \quad (191)$$

The cross-derivative condition $U_{xt} = U_{tx}$ is solved for S_t

$$S_t = 4\lambda(t)C_x - 2\lambda'(t). \quad (192)$$

Substituting U_x, U_t, S_t and C_{xxx} taken from (62) in E_5 , one obtains the condition

$$E_5 = 2\alpha\lambda(t)\lambda'(t) = 0, \quad (193)$$

which introduces the spectral parameter as the arbitrary constant λ .

The solution for (U_x, U_t) is

$$U_x - \beta = (\alpha/4)(S + 2\lambda), \quad U_t - \gamma = \alpha\lambda C, \quad (194)$$

and the elimination of U defines the SME

$$\frac{S_t}{C_x} - 4\lambda = 0. \quad (195)$$

The solution for (S, C) is

$$S = (4/\alpha)(U_x - \beta) - 2\lambda, \quad C = (U_t - \gamma)/(\alpha\lambda), \quad (196)$$

and its cross-derivative condition

$$X \equiv E(U)/(\alpha\lambda) = 0 \quad (197)$$

creates on the field U the only constraint that U satisfy the AKNS PDE.

The BT is the result of the substitution $\chi^{-1} = (u - U)/\alpha$ in (64)–(65).

The KdV equation

The Korteweg-de Vries equation for u (56) is defined in conservative form, so it is cheaper to process the potential form

$$E(v) \equiv bv_t + v_{xxx} - (3/a)v_x^2 + F(t) = 0, \quad v = u_x. \quad (198)$$

Its unique family is

$$p = -1, \quad q = -4, \quad v_0 = -2a, \quad \text{indices } (-1, 1, 6), \quad \mathcal{D} = -2a\partial_x. \quad (199)$$

With the assumption

$$v = V + \mathcal{D} \text{Log } \tau, \quad E(v) = 0, \quad (200)$$

for the DT, the choice of the second-order scalar (111)–(112) for ψ , the link (146) between τ and ψ , one generates the three determining equations

$$E_2 \equiv -2a(bC + 2S) - 12V_x = 0, \quad (201)$$

$$E_3 \equiv 2a(bC - S)_x = 0, \quad (202)$$

$$E_4 \equiv E(V) + \frac{S}{2}E_2 - \frac{1}{2}E_{3,x} = 0. \quad (203)$$

After one integration of E_3 , the system (E_2, E_3) is solved for (S, C)

$$S = -2\lambda(t) - (2/a)V_x, \quad bC = 4\lambda(t) - (2/a)V_x, \quad (204)$$

in which $\lambda(t)$ is an arbitrary integration function. Then E_4 , as seen from its above written compacted expression, expresses that V satisfies the PDE. Last, the cross-derivative condition (62)

$$X \equiv -2\lambda'(t) - 2(E(V))_x/(ab) = 0 \quad (205)$$

introduces the spectral parameter as an arbitrary complex constant and proves that a Lax pair has been obtained for the conservative (not the potential) equation. This Lax pair can be written, at the reader's taste, either in the scalar representation (111)–(112), with $U = V_x$,

$$L_1 \equiv \partial_x^2 - U/a - \lambda, \quad (206)$$

$$L_2 \equiv b\partial_t + (4\lambda - 2U/a)\partial_x + U_x/a, \quad (207)$$

$$a[L_1, L_2] = bU_t + U_{xxx} - (6/a)UU_x, \quad (208)$$

or in the zero-curvature representation (115)–(116)

$$L = \begin{pmatrix} 0 & 1 \\ U/a + \lambda & 0 \end{pmatrix}, \quad (209)$$

$$M = b^{-1} \begin{pmatrix} -U_x/a & 2U/a - 4\lambda \\ -U_{xx}/a + 2(U/a + \lambda)(U/a - 2\lambda) & U_x/a \end{pmatrix}, \quad (210)$$

or in the Riccati representation for $\omega = \chi^{-1}$ (see (64)–(65) and (69))

$$\omega_x = -\frac{S}{2} - \omega^2 = \left(\frac{U}{a} + \lambda\right) - \omega^2, \quad (211)$$

$$\omega_t = (-C\omega + C_x/2)_x = b^{-1}((2U/a - 4\lambda)\omega - U_x/a)_x. \quad (212)$$

This last representation is by far the best one, for it allows one to deduce immediately two quite important informations, namely the auto-Bäcklund transformation of KdV and the hetero-Bäcklund transformation between KdV and mKdV. Firstly, the substitution of the inverse relation of (200)

$$\omega = (v - V)/(-2a) \quad (213)$$

in (211)–(212) provides the auto-BT for the conservative form of KdV

$$a(v + V)_x = -2a^2\lambda + (v - V)^2/2, \quad (214)$$

$$a(b(v + V)_t - 2F'(t)) = -(v - V)(v - V)_{xx} + 2(V_x^2 + v_x V_x + v_x^2), \quad (215)$$

after suitable differential consequences of the x -part have been added to the t -part in order to suppress λ and cubic terms in (215).

Secondly, the elimination of U between (211)–(212) leads to the mKdV equation (404) for w , with the identification $w = \alpha\omega, \nu = \lambda$; since conversely the elimination of ω leads to the KdV equation for U , the system (211)–(212) also represents the hetero-BT between KdV and mKdV ([71] Eq. (5.16), [112]).

As to the SME, it results from the elimination of V between (E_2, E_3, E_4)

$$bC - S - 6\lambda = 0. \quad (216)$$

Most of these results for KdV were found in the original paper of WTC [119].

Remark. The transformation (211) between $w = \alpha\omega$ and U is often called a *Miura transformation*, but it is really just one half of the hetero-BT. The advantage of the hetero-BT is that it is invertible, while the Miura transformation as defined in the previous sentence is not.

7.2 Integrable equations with a third order Lax pair

Let us process a few PDEs which possess a third order Lax pair, and let us first perform their one-family truncation with the (wrong) assumption of a second order Lax pair, because this often provides interesting results.

The Boussinesq equation

The Boussinesq equation (Bq) is often defined in a two-component evolution form [124]

$$\text{sBq}(u, r) \equiv \begin{cases} u_t - r_x = 0, & (\alpha, \beta, \varepsilon) \text{ constant,} \\ r_t + \varepsilon^2((u + \alpha)^2 + (\beta^2/3)u_{xx})_x = 0. \end{cases} \quad (217)$$

Let us consider its one-component “potential” form

$$\text{pBq}(v) \equiv v_{tt} + \varepsilon^2((v_x + \alpha)^2 + (\beta^2/3)v_{xxx})_x = 0, \quad u = v_x, \quad r = v_t. \quad (218)$$

Equation (218) has only one family of movable singularities

$$p = -1, \quad q = -5, \quad \text{indices } (-1, 1, 4, 6), \quad \mathcal{D} = 2\beta^2\partial_x, \quad (219)$$

and it passes the Painlevé test [117]. Since (218) is a conservation law, the computations can be reduced by considering the “second potential Bq” equation

$$\text{ppBq}(w) \equiv w_{tt} + \varepsilon^2((w_{xx} + \alpha)^2 + (\beta^2/3)w_{xxxx}) = 0, \quad u = v_x = w_{xx}, \quad (220)$$

whose single family is of the logarithmic type $w \sim 2\beta^2 \text{Log } \chi$

$$p = 0^-, q = -4, \text{ indices}(-1, 0, 1, 6), \mathcal{D} = 2\beta^2. \quad (221)$$

Let us assume for the would-be DT the relation

$$w = 2\beta^2 \text{Log } \tau + W, \text{ ppBq}(w) = 0, \quad (222)$$

and for the link between τ and ψ the identity (146).

Let us first assume that ψ satisfies the second-order scalar Lax pair (111)–(112). This is equivalent to the usual WTC truncation in the invariant formalism [19]

$$\text{ppBq}(w) \equiv \sum_{j=0}^4 E_j \chi^{j-4} = 0, \quad (223)$$

and this generates the three determining equations

$$E_2 \equiv (4/3)\beta^2 \varepsilon^2 S - 2C^2 - 4\varepsilon^2(W_{xx} + \alpha) = 0, \quad (224)$$

$$E_3 \equiv -2(C_t - CC_x - (\beta^2 \varepsilon^2/3)S_x) = 0, \quad (225)$$

$$E_4 \equiv (SE_2 - E_{3,x})/2 + C_x^2 + \beta^2 \text{ppBq}(W) = 0. \quad (226)$$

From the last equation $E_4 = 0$, the desired solution $\text{ppBq}(W) = 0$ cannot be generic, so this second-order assumption fails to provide the auto-BT. However, it does provide another information, namely a hetero-BT between the Boussinesq PDE and another PDE. Indeed, under the natural parametric representation of E_3 (which, by the way, would be the SME if the second order were the correct one),

$$S = 3z_t - 3(\beta\varepsilon)^2 z_x^2/2, \quad C = (\beta\varepsilon)^2 z_x, \quad (227)$$

the field z , by the cross-derivative condition (62), satisfies the modified Boussinesq equation [61]

$$\text{MBq}(z) \equiv z_{tt} + ((\beta\varepsilon)^2/3)z_{xxxx} + 2(\beta\varepsilon)^2 z_t z_{xx} - 2(\beta\varepsilon)^4 z_x^2 z_{xx} = 0. \quad (228)$$

Just like for the KdV equation (Section 7.1), this leads, after a short computation left to the reader, to the hetero-BT between the Boussinesq and the modified Boussinesq equations.

Going to third order, the assumption (222) and (146), with ψ solution of the scalar Lax pair (117)–(118), generates

$$\text{ppBq}(w) \equiv \sum_{k=0}^2 \sum_{l=0}^2 E_{k,l} Y_1^k Y_2^l, \quad k+l \leq 2. \quad (229)$$

These six determining equations $E_{k,l} = 0$, plus the three cross-derivative conditions $X_j = 0, j = 0, 1, 2$, are solved as follows in the Gel'fand-Dikii case $f = 0$: [87,90]

$$\begin{aligned}
 E_{02} &\equiv (\beta\varepsilon)^2 - c^2 = 0 && \Rightarrow c = \beta\varepsilon, \\
 E_{11} &\equiv d = 0 && \Rightarrow d = 0, \\
 E_{20} &\equiv 3(V_x + \alpha) + 2\beta^2 a = 0 && \Rightarrow a = -3(V_x + \alpha)/(2\beta^2), \\
 E_{10} &\equiv \varepsilon V_{xx} - \beta e_x = 0 && \Rightarrow e_x = \beta^{-1}\varepsilon V_{xx}, \\
 X_1 &\equiv 3V_{xt} + 3\beta\varepsilon V_{xxx} + 4\beta^3\varepsilon b_x = 0 && \Rightarrow b = g(t) - 3(\beta^{-2}V_{xx} + \beta^{-3}\varepsilon^{-1}V_t)/4, \\
 X_0 &\equiv (3/(4\varepsilon\beta^2))\text{pBq}(V) = 0 && \Rightarrow V \text{ satisfies the PDE (218),} \\
 E_{00} &\equiv 2\beta^2 g'(t) = 0 && \Rightarrow g(t) = \lambda,
 \end{aligned} \tag{230}$$

in which λ is an arbitrary constant. The coefficients a, b, c, d, e are

$$\begin{aligned}
 a &= -(3/2)\beta^{-2}(V_x + \alpha), \quad b = \lambda - (3/4)\beta^{-2}V_{xx} - (3/4)\beta^{-3}\varepsilon^{-1}V_t, \\
 c &= \beta\varepsilon, \quad d = 0, \quad e = \beta^{-1}\varepsilon(V_x + \alpha),
 \end{aligned} \tag{231}$$

$$X_0 = (3/(4\varepsilon\beta^2))\text{pBq}(V), \quad X_1 = 0, \quad X_2 = 0, \tag{232}$$

and they define a third-order Lax pair of the potential Boussinesq equation (218) [123,124,82].

The BT is just (152)–(153) or equivalently, after substitution of $Y_1 = (v - V)/(2\beta^2)$,

$$\begin{aligned}
 (v - V)_{xx} + 3\beta^{-1}\varepsilon^{-1}(v + V)_t + 3\beta^{-2}(v - V)((v + V)_x + 2\alpha) \\
 + \beta^{-4}(v - V)^3 - 8\beta^2\lambda = 0,
 \end{aligned} \tag{233}$$

$$(v + V)_{xx} - \beta^{-1}\varepsilon^{-1}(v - V)_t + \beta^{-2}(v - V)(v - V)_x = 0. \tag{234}$$

The Hirota-Satsuma equation

Defined as [60]

$$\text{HS}(w) \equiv [w_{xxt} + (6/a)w_x w_t]_x = 0, \quad a \neq 0, \tag{235}$$

it is better processed on its potential form

$$\text{pHS}(w) \equiv w_{xxt} + (6/a)w_x w_t + F(t) = 0, \quad a \neq 0. \tag{236}$$

The second order assumption (111)–(112) generates the three determining equations

$$\begin{aligned}
 E_2 &\equiv -2aSC - 6W_t + 6CW_x = 0, \\
 E_3 &\equiv aS_t + 2aSC_x - 6C_x W_x = 0, \\
 E_4 &\equiv \text{pHS}(W) - a(S^2 C + S_{xt}/2 + SC_{xx}) - 3SW_t + 3(SC + C_{xx})W_x = 0.
 \end{aligned} \tag{237}$$

In the generic case $C_x \neq 0$, their general solution is unknown, in particular we have not succeeded to perform the elimination of (S, C) to find the constraint(s) satisfied by W . It is easy to eliminate W but this gives rise to two

equations for (S, C)

$$6W_x = 2aS + a\frac{S_t}{C_x}, \quad 6W_t = a\frac{CS_t}{C_x}, \quad (238)$$

$$M_{23} \equiv \left(\frac{CS_t}{C_x}\right)_x - \left(2S + \frac{S_t}{C_x}\right)_t = 0, \quad (239)$$

$$M_4 \equiv 1 - 6a^{-1}F(t) - C_x^{-1}(4CSS_t + CS_{xxt} + 2C_{xx}S_t) - C_x^{-2}(2CS_t^2 + C^2S_tS_x - 2CC_{xx}S_{xt}) - 2C_x^{-3}CC_{xx}^2S_t = 0, \quad (240)$$

and their possible functional dependence is unsettled. Anyhow, the field W cannot be a second solution of (235) [86].

The third order assumption (117)–(118), with the link (146) and the truncated expansion

$$w = W + a\partial_x \text{Log } \tau, \quad (241)$$

generates seven determining equations (150). They are easily solved [86] and their unique solution defines the Lax pair (268)–(269), with W a second solution of (236).

The Tzitzéica equation

The equation is defined by (168), in the case $(a_1 = 0, a_2 \neq 0)$. It possesses two families, the first one defined by (169), the second one by

$$e^{-u} \sim \sqrt{(1/a_2)\varphi_x\varphi_t}(\varphi - \varphi_0)^{-1}, \quad \text{indices } (-1, 2). \quad (242)$$

These two families are not opposite, but the second family is irrelevant because the Tzitzéica equation has a one-to-one correspondence [59] with a one-family equation, namely the potential form (236) of the Hirota-Satsuma PDE in the particular case $F(t) = 0$. This correspondence is obtained by the elimination of a_2 in equation (168)

$$\left(F(t) = 0, e^u = \frac{2}{a\alpha}w_t\right) \implies \left(e^{-2u}(e^{2u}\text{Tzi}(u))_x = \left(\frac{\text{pHS}(w)}{w_t}\right)_t\right). \quad (243)$$

The irrelevance of the second family is confirmed by the negative result of Weiss [118,19] obtained when performing a truncation on e^{-u} .

All the truncations will accordingly take the same form (170) as for the Liouville equation, which implies that τ is an object invariant under the permutation (131). Depending on the Lax pair assumption, the link between τ and ψ will be either the identity (case of a scalar ψ invariant under the permutation (131)) or not (if the scalar ψ is not invariant, e.g. because the Lax pair itself is not invariant), as detailed below.

Let us first assume a second order Lax pair. To the author's knowledge, one cannot define a scalar ψ , linked to such a Lax pair, which, like τ , would

be invariant under (131). This is probably the reason why the assumption $\tau = \psi$ with ψ solution of the noninvariant Lax pair (111)–(112) generates so intricate determining equations that their general solution has not yet been obtained [88]; these equations are however consistent in the sense that one easily finds the particular exact solution

$$\alpha e^u = 2c\wp(x - ct - x_1, g_2, A + \frac{a_2\alpha^2}{8c^3}) - 2c\wp(x + ct - x_2, g_2, A - \frac{a_2\alpha^2}{8c^3}), \quad (244)$$

depending on five arbitrary constants (x_1, x_2, c, g_2, A) and representing the superposition of two traveling waves of opposite velocities.

From this second-order WTC truncation, and with appropriate assumptions, one can also find a particular solution which represents a binary Darboux transformation [108].

Let us now turn to the third order assumption. One can postulate either a Lax pair invariant under (131), such as the matrix pair (132)–(133), or a noninvariant Lax pair such as the scalar pair (117)–(118). In the first case, one must assume the identity link $\tau = \psi$, while in the second case the assumed link must be noninvariant. Both assumptions lead to a success [37]. Let us detail here the invariant assumption, i.e. *a priori* the simpler one.

The truncation is defined by (170), the link (146), and the matrix Lax pair (132)–(133)

$$E(u) \equiv \sum_{k=0}^3 \sum_{l=0}^{3-k} E_{kl}(f_j, g_j, h_j, U) Y_1^k Y_2^l, \quad \forall k, l : E_{kl} = 0, \quad (245)$$

in which (Y_1, Y_2) are the two components of the projective Riccati pseudopotential (135)–(138) equivalent to the Lax pair. To these ten determining equations in U and the nine unknown coefficients, one must add the six cross-derivative conditions $X_j = 0$ (139)–(140).

During their resolution, one first proves that the product $f_2 h_1$ cannot vanish (otherwise a_2 would be zero). This makes the sixteen equations algebraically independent and equivalent to the fifteen differential relations

$$f_{j,t}, g_{j,x}, g_{j,t}, h_{j,x}, g_{j,xt} = P(\{f_k, g_k, h_k\}, k = 1, 2, 3), \quad j = 1, 2, 3, \quad (246)$$

with P polynomials whose coefficients depend on U, U_x, U_t, U_{xt} , plus the single algebraic relation

$$E_{00} \equiv a_2 - \frac{4}{\alpha^2} (g_3 + g_1 g_2 + (\alpha/2)e^U)^2 = 0. \quad (247)$$

They are solved successively as [equations are referenced as in (246)–(247)]

$$\begin{aligned}
g_{3,xt} - (g_{3,x})_t &: E(U) = 0, \\
g_{1,x} - g_{2,t} &: \exists g_0(x, t) : g_1 = g_{0,t}, \quad g_2 = g_{0,x}, \\
g_{2,t} &: g_3 = -\alpha e^U - g_{0,x}g_{0,t} - g_{0,xt}, \\
E_{00} &: \exists f_0(x, t) \neq 0 : f_2 = \sqrt{a_2}W^{-1}f_0, \quad h_1 = \sqrt{a_2}W^{-1}f_0^{-1}, \\
&\quad \text{notation } W = e^U + (2/\alpha)g_{0,xt}, \\
g_{2,x} &: f_3 = -\sqrt{a_2}W^{-1}f_0g_{0,t} - f_1g_{0,x} - g_{0,x}^2 + g_{0,xx}, \\
g_{3,x} &: f_1 = W_x/W + 2g_{0,x}, \\
f_{2,t} &: h_2 = W_t/W + 2g_{0,t} - f_{0,t}/f_0, \\
h_{1,x} &: f_{0,x} = 0, \\
g_{1,t} &: h_3 = g_{0,t}(f_{0,t}/f_0 - W_t/W - g_{0,t}) + g_{0,tt} - \sqrt{a_2}W^{-1}g_{0,x}/f_0, \\
g_{3,t} &: f_{0,t} = 0, \\
h_{2,x} &: g_{0,xt} = 0.
\end{aligned} \tag{248}$$

The irrelevant arbitrary function g_0 reflects the freedom in the definition (170) of τ and can be absorbed by redefining τ as τe^{-g_0} . Thus the solution is unique : the field U must satisfy the Tzitzéica PDE, and f_0 is an arbitrary nonzero complex constant λ . Accordingly, one has obtained a Lax pair and a Darboux transformation. The equivalent projective Riccati representation of the matrix Lax pair is

$$Y_{1,x} = -Y_1^2 + U_x Y_1 + \sqrt{a_2} \lambda e^{-U} Y_2, \tag{249}$$

$$Y_{2,x} = -Y_1 Y_2 - \alpha e^U, \tag{250}$$

$$Y_{1,t} = -Y_1 Y_2 - \alpha e^U, \tag{251}$$

$$Y_{2,t} = -Y_2^2 + U_t Y_2 + \sqrt{a_2} \lambda^{-1} e^{-U} Y_1, \tag{252}$$

with cross-derivative conditions proportional to the Tzitzéica equation

$$(Y_{1,x})_t - (Y_{1,t})_x = Y_1 E(U), \quad (Y_{2,x})_t - (Y_{2,t})_x = Y_2 E(U). \tag{253}$$

This Lax pair is the rewriting in matrix form of the scalar triplet given by Tzitzéica [110]

$$-\tau_{xx} + U_x \tau_x + \sqrt{a_2} \lambda e^{-U} \tau_t = 0, \tag{254}$$

$$-\tau_{tt} + U_t \tau_t + \sqrt{a_2} \lambda^{-1} e^{-U} \tau_x = 0, \tag{255}$$

$$-\tau_{xt} - \alpha e^U \tau = 0. \tag{256}$$

The Lax pair admits by construction the involution [111,50]

$$(\tau, e^U, \lambda) \rightarrow \left(\frac{1}{\tau}, -e^U - \frac{2}{\alpha} \frac{\tau_x \tau_t}{\tau^2}, -\lambda \right), \tag{257}$$

equivalent to

$$(\tau, e^U, \lambda) \rightarrow (1/\tau, e^U + \mathcal{D} \text{Log } \tau, -\lambda), \tag{258}$$

which defines another, equivalent, writing of the Darboux transformation

$$e^u = -e^U - \frac{2}{\alpha} \frac{\tau_x \tau_t}{\tau^2}. \quad (259)$$

Remark. Knowing these results, one can also write this DT [122,3] as a difference of the two fields $u - U$ in terms of the two components of a projective Riccati pseudopotential

$$u = U + \text{Log}(-2\lambda^2 y_1 y_2 - 1), \quad y_j = \alpha^{-1/2} \lambda^{-1} e^{-U/2} Y_j, \quad (260)$$

in a quite similar manner to the DT of Liouville and sine-Gordon (392). However, the field u is multivalued.

In order to find the BT, one must now eliminate one of the two equivalent projective components, and this defines two possible, different, eliminations.

In the first elimination, one takes Y_2 from (249) and substitutes it into the three remaining equations, which results in

$$Y_2 = (Y_{1,x} + Y_1^2 - U_x Y_1) e^U / (\sqrt{a_2} \lambda), \quad (261)$$

$$\text{ODE} \equiv Y_{1,xx} + 3Y_1 Y_{1,x} + Y_1^3 - e^{-U} (e^U)_{xx} Y_1 + \alpha \sqrt{a_2} \lambda = 0, \quad (262)$$

$$\text{PDE} \equiv Y_{1,t} + e^U ((Y_1 Y_{1,x} + Y_1^3) - Y_1^2 U_x) / (\sqrt{a_2} \lambda) + \alpha e^U = 0, \quad (263)$$

$$(252) \equiv -Y_1 E(U) - \frac{e^U Y_1}{\sqrt{a_2} \lambda} \text{ODE} + (2Y_1 - U_x + \partial_x) \text{PDE} = 0, \quad (264)$$

$$[\text{ODE}, \text{PDE}] = (Y_{1,xx})_t - (Y_{1,t})_{xx} = Y_1 (e^{2U} E(U))_x. \quad (265)$$

Only two of them are functionally independent, as shown by relation (264), but the commutator (265) of equations (262)–(263) shows that this elimination fails to generate the auto-BT of Tzitzéica equation.

However, it does provide another result, which we now derive. The ODE (262) belongs to the classification of Gambier – this is the number 5, see Section 5.6 –, it is linearizable by the transformation $Y_1 = \partial_x \text{Log} \psi$ into a third-order linear ODE, with the relation $\tau = \psi$ between the two functions. This transformation also linearizes the PDE (263), and the resulting linear system

$$\tau_{xxx} - (U_{xx} + U_x^2) \tau_x + \sqrt{a_2} \alpha \lambda \tau = 0, \quad (266)$$

$$-\sqrt{a_2} \lambda \tau_t + e^U \tau_{xx} - U_x e^U \tau_x = 0, \quad (267)$$

which cannot be a scalar Lax pair of the Tzitzéica equation, is, in fact, the scalar Lax pair of the Hirota-Satsuma equation (235), see Section 7.2,

$$\tau_{xxx} - (6/a) w_x \tau_x + \Lambda \tau = 0, \quad (268)$$

$$\Lambda \tau_t - (2/a) w_t \tau_{xx} + (2/a) w_{xt} \tau_x = 0, \quad (269)$$

under the change of variables (243).

In the second elimination, one takes Y_1 from (250) and substitutes it into the three remaining equations

$$Y_1 = -(Y_{2,x} + \alpha e^U)/Y_2, \quad (270)$$

$$\begin{aligned} \text{ODE} \equiv & Y_2 Y_{2,xx} - 2Y_{2,x}^2 - (U_x Y_2 + 3\alpha e^U) Y_{2,x} \\ & + \sqrt{a_2} \lambda e^{-U} Y_2^3 - \alpha^2 e^{2U} = 0, \end{aligned} \quad (271)$$

$$\text{PDE} \equiv Y_2 Y_{2,t} + Y_2^3 - U_t Y_2^2 + \sqrt{a_2} \lambda^{-1} (\alpha + e^{-U} Y_{2,x}) = 0, \quad (272)$$

$$\begin{aligned} (251) \equiv & E(U) + (\partial_x - \alpha e^U Y_2^{-1}) \text{PDE} \\ & - \sqrt{a_2} \lambda^{-1} e^{-U} Y_2^{-2} \text{ODE} = 0, \end{aligned} \quad (273)$$

$$\begin{aligned} [\text{ODE}, \text{PDE}] &= (Y_{2,xx})_t - (Y_{2,t})_{xx} \\ &= (3\alpha e^U + U_x Y_2 + 3Y_{2,x} - Y_2 \partial_x) E(U). \end{aligned} \quad (274)$$

Only two of them are functionally independent, as shown by the relation (273), and the vanishing of the commutator (274) of equations (271)–(272) is equivalent to the vanishing of the Tzitzéica equation for U . This elimination therefore generates the auto-BT of Tzitzéica equation, by the substitution

$$Y_2 = (\alpha/2) \int (e^u - e^U) dx \quad (275)$$

into (271)–(272).

The ODE part (271) of the BT is equivalently written as [91]

$$\frac{w_{xx}}{w_x} - \frac{W_{xx}}{W_x} - 2 \frac{w_x + W_x}{w - W} + \alpha \sqrt{a_2} \lambda \frac{(w - W)^2}{2w_x W_x} = 0, \quad (276)$$

with the notation $Y_2 = (\alpha/2)(w - W)$, $e^u = w_x$, $e^U = W_x$.

The nonlinear ODE (271) again belongs to the equivalence class of the fifth Gambier equation (G5), see section 5.6, and its linearization

$$Y_2^{-1} = -\alpha^{-1} e^{-U} \partial_x \text{Log}(e^U \psi) \quad (277)$$

transforms the two equations (271)–(272) into the third-order scalar Lax pair of the Gel'fand and Dikii type (i.e. $f = 0$ in (117)–(118))

$$\mathcal{L}\psi \equiv \psi_{xxx} + (2U_{xx} - U_x^2)\psi_x + ((2U_{xx} - U_x^2)_x/2 + \sqrt{a_2}\alpha\lambda)\psi = 0, \quad (278)$$

$$\begin{aligned} \mathcal{M}\psi \equiv & \psi_t + \sqrt{a_2}(\alpha\lambda)^{-1} e^{-2U} (\psi_{xx} + U_x \psi_x + U_{xx} \psi) \\ & + (U_t + \int (\alpha e^U + a_2 e^{-2U}) dx) \psi = 0, \end{aligned} \quad (279)$$

$$\begin{aligned} [\mathcal{L}, \mathcal{M}] &= 3E\partial_x^2 + (2(e^U)_x E + E_x)\partial_x \\ &+ (e^U)_x E_x + (3U_{xx} - U_x^2)E, \quad E = E(U). \end{aligned} \quad (280)$$

Thus, the noninvariant (under (131)) link between τ and ψ that one would have had to postulate if one had chosen the scalar Lax pair (117)–(118) is a

posteriori provided by the linearizing formula (277) and the Riccati equation (251), this is the invertible transformation

$$e^U \tau = (e^U \psi)_x, \quad e^U \psi = -\alpha^{-1} \tau_t, \quad (281)$$

and it clearly breaks the invariance under (131).

The Sawada-Kotera and Kaup-Kupershmidt equations

Because of their duality [65,115], it is convenient to introduce simultaneously the Sawada-Kotera equation (SK) and the Kaup-Kupershmidt equation (KK). These are defined as

$$\text{SK}(u) \equiv \beta u_t + \left(u_{xxxx} + \frac{30}{\alpha} u u_{xx} + \frac{60}{\alpha^2} u^3 \right)_x = 0, \quad (282)$$

$$\text{pSK}(v) \equiv \beta v_t + v_{xxxx} + \frac{30}{\alpha} v_x v_{xxx} + \frac{60}{\alpha^2} v_x^3 = 0, \quad (283)$$

$$\text{KK}(u) \equiv \beta u_t + \left(u_{xxxx} + \frac{30}{\alpha} u u_{xx} + \frac{45}{2\alpha} u_x^2 + \frac{60}{\alpha^2} u^3 \right)_x = 0, \quad (284)$$

$$\text{pKK}(v) \equiv \beta v_t + v_{xxxx} + \frac{30}{\alpha} v_x v_{xxx} + \frac{45}{2\alpha} v_{xx}^2 + \frac{60}{\alpha^2} v_x^3 = 0, \quad (285)$$

in which u denotes the conservative field and v the potential one, with $u = v_x$.

Both equations have the Painlevé property [115]. Each of them has two families [115]

$$\text{pSK, F1} : p = -1, \quad v_0 = \alpha, \quad \text{indices } -1, 1, 2, 3, 10, \quad (286)$$

$$\text{pSK, F2} : p = -1, \quad v_0 = 2\alpha, \quad \text{indices } -2, -1, 1, 5, 12, \quad (287)$$

$$\text{pKK, F1} : p = -1, \quad v_0 = \alpha/2, \quad \text{indices } -1, 1, 3, 5, 7, \quad (288)$$

$$\text{pKK, F2} : p = -1, \quad v_0 = 4\alpha, \quad \text{indices } -7, -1, 1, 10, 12. \quad (289)$$

The singular part operator \mathcal{D} attached to a given family is $\mathcal{D} = v_0 \partial_x$. The two families have residues which are not opposite, but fortunately each potential equation possesses in its hierarchy a “minus-one” equation [120]

$$\text{pSK}_{-1} : v_{xxt} + \frac{6}{\alpha} v_x v_t = 0, \quad (290)$$

$$\text{pKK}_{-1} : v_t v_{xxt} - \frac{3}{4} v_{xt}^2 + \frac{6}{\alpha} v_x v_t^2 = 0, \quad (291)$$

which has only one family (the first one is nothing else than the Hirota-Satsuma PDE, already processed in Section 7.2). The equations SK and KK are therefore to be considered as possessing the single family F1, Eqs. (286) and (288).

Let us assume the one-family DT (200) and, successively, the second-order scalar Lax pair (111)–(112), then the third-order scalar one (117)–(118) with

the Gel'fand-Dikii simplification $f = 0$. As to the link between τ and ψ , at second order this is the identity, while at third order it can be, as outlined in Section 5.6 and detailed in Musette lecture [85], either the linearizing transformation of the fifth Gambier equation or that of the twenty-fifth Gambier equation.

Therefore, at the fourth step of the singular manifold method, for each PDE, one has only three possibilities to examine : order two and Riccati, order three and (G5), order three and (G25). This is done in the next two Sections.

The Sawada-Kotera equation

First truncation (order two and Riccati). The one-family truncation (147) with $\tau = \psi$ generates the three equations [20]

$$E_4 \equiv \beta C - 4S^2 + 9S_{xx} + 60SV_x/\alpha - 180(V_x/\alpha)^2 - 30V_{xxx}/\alpha = 0, \quad (292)$$

$$E_5 \equiv -\beta C_x - 2SS_x + S_{xxx} + 30S_x V_x/\alpha = 0, \quad (293)$$

$$E_6 \equiv \text{pSK}(V) + (SE_4 - E_{5,x})/2 + 5S_x(3V_{xx}/\alpha - S_x/2) = 0. \quad (294)$$

These equations possess two solutions [20], a nongeneric one $S_x = 0$

$$\begin{aligned} S &= -k^2/2, \quad C = c + c_0, \quad c = k^4/\beta + 2c_0/3, \\ V/\alpha &= \zeta(x - (c - c_0)t, k^4/12 + \beta c_0/9, g_3) - k^2x/12 \\ &\quad + ((5(k^4 + \beta c_0)k^2/36 - 12g_3)t)/\beta, \end{aligned} \quad (295)$$

in which ζ is the Weierstrass function and (k, c_0, g_3) are arbitrary constants, and a generic one $S_x \neq 0$ defined by the four equations

$$V_x = \alpha(\beta C_x + 2SS_x - S_{xxx})/(30S_x), \quad (296)$$

$$V_t = \dots \quad (297)$$

$$M_1 \equiv (G_x/S_x)_{xx} - G - S_x^2 G_x^2/5 + 2SG_x/S_x = 0, \quad (298)$$

$$M_2 \equiv 29 \text{ terms} = 0, \text{ also vanishing if } G = 0, \quad (299)$$

in which G is defined by

$$G \equiv S_{xx} + 4S^2 - \beta C. \quad (300)$$

The general solution (S, C) of the system $M_1 = 0, M_2 = 0$ has not yet been obtained, for the elimination of S or C is difficult. This difficulty reflects the fact that second order is not the correct order. Nevertheless, these complicated equations admit the very simple *particular* solution [115] (it would be interesting to prove that this is the *general* solution),

$$G = 0, \quad V_x = \alpha S/3, \quad \text{pKK}(V) = 0, \quad (301)$$

so the field V in the DT assumption (200) satisfies a different PDE, namely the potential KK equation. This defines a hetero-BT between the conservative forms of SK and KK [46,59,58]

$$\begin{aligned} \alpha(v + V/2)_x + (v - V)^2 = 0, \quad \beta(v + V/2)_t + \dots = 0, \\ \text{pSK}(v) = 0, \quad \text{pKK}(V) = 0 \end{aligned} \quad (302)$$

(see Ref. [58] for the exact expression of the t -part).

With $G = 0$, the linear system (111)–(112) is a degenerate Lax pair for KK, since it lacks a spectral parameter.

Still when (301) holds, the field χ^{-1} satisfies a fifth order PDE, the Fordy-Gibbons equation, and the explicit writing of its hetero-BT with the SK equation is left to the reader.

Second truncation (order three and (G5)). This assumption creates no *a priori* constraint on the coefficients (a, b) of the spectral problem (117), and the linearizing transformation of (G5) is just the identity $\tau = \psi$. This generates six determining equations (150). The process is successful [115,118,22] and V is found to be a second solution of pSK (notation $U = V_x$ as usual)

$$b = \lambda, \quad (303)$$

$$a = -6U/\alpha, \quad (304)$$

$$L_1 = \partial_x^3 + 6\frac{U}{\alpha}\partial_x - \lambda, \quad (305)$$

$$L_2 = \beta\partial_t + \left(18\frac{U_x}{\alpha} - 9\lambda\right)\partial_x^2 + \left(36\frac{U^2}{\alpha^2} - 6\frac{U_{xx}}{\alpha}\right)\partial_x - 36\lambda\frac{U}{\alpha}, \quad (306)$$

$$[L_1, L_2] = 6\beta^{-1}\alpha^{-1}\text{SK}(U). \quad (307)$$

This is the Lax pair given by Satsuma and Kaup [106].

The BT results from the elimination of Y_2 , which provides Eqs. (152)–(153) for $Y_1 = Y$,

$$Y_{xx} + 3YY_x + Y^3 + 6(U/\alpha)Y - \lambda = 0, \quad (308)$$

$$\beta Y_t - 9[(\lambda - 2U_x/\alpha)(Y_x + Y^2) \quad (309)$$

$$+ 4(\lambda U/\alpha - (U/\alpha)^2 Y) + (2/3)(U_{xx}/\alpha)Y]_x = 0,$$

$$\beta((Y_{xx})_t - (Y_t)_{xx})/Y = -(6/\alpha)\text{SK}(U), \quad (310)$$

followed by the substitution $Y = (v - V)/\alpha$,

$$(v - V)_{xx}/\alpha + 3(v - V)(v + V)_x/\alpha^2 + (v - V)^3/\alpha^3 - \lambda = 0, \quad (311)$$

$$\begin{aligned} \beta(v - V)_t/\alpha - (3/2)[(v - V)_{xxx}/\alpha \\ + (5(v - V)(v + V)_{xxx} + 15(v + V)_x(v - V)_{xx})/\alpha^2 \\ + (15(v - V)^2(v - V)_{xx} + 30(v - V)(v + V)_x^2)/\alpha^3 \\ + 30(v - V)^3(v + V)_x/\alpha^4 + 6(v + V)^5/\alpha^5]_x = 0, \end{aligned} \quad (312)$$

a result due to Satsuma and Kaup [106].

The Kaup-Kupershmidt equation

First truncation (order two and Riccati). The one-family truncation (147) with $\tau = \psi$ generates the three equations [90]

$$E_2 \equiv 15(S/4 - 3V_x/\alpha) = 0, \quad (313)$$

$$E_4 \equiv \beta C/2 + 7S^2/4 + 3S_{xx}/4 - 15(SV_x + V_{xxx})/\alpha - 90(V_x/\alpha)^2 = 0, \quad (314)$$

$$E_6 \equiv (S/2)E_4/\alpha + (4\beta(CS + C_{xx}) - S^3 + (21/2)S_x^2 + 14SS_{xx} - 4S_{xxx})/16 + (15/4)(3S^2 - 2S_{xx})V_x/\alpha - 45S(V_x/\alpha)^2 + \text{pKK}(V) = 0. \quad (315)$$

As opposed to the (difficult) SK case, these equations are easy to solve and possess the unique solution [115]

$$V_x = \alpha S/12, \text{ pSK}(V) = 0, S_{xx} + S^2/4 - \beta C = 0. \quad (316)$$

This is a strong indication that the particular solution (301) of (296)–(299) should be the general one. One again recovers, by a nice duality, the hetero-BT between KK and SK.

Second truncation (order three and (G5)). This generates thirteen determining equations (150). This truncation fails and provides no solution at all (one determining equation is $E_{2,2} \equiv 45/8 = 0$), not even the one-soliton solution. Indeed, the one-soliton solution of Kaup corresponds to constant coefficients for the scalar Lax pair (117)–(118) with $f = 0$, and, with the above procedure, the only way to obtain it [31,101] is to enforce the two first integrals K_1 and K_2 which result from the zero value of b ,

$$K_1 = \psi_{xx} - a\psi, K_2 = \psi_x^2 - a\psi^2 - 2(\psi_{xx} - a\psi)\psi. \quad (317)$$

Third truncation (order three and (G25)). This assumption implies, among the coefficients (a, b) of the spectral problem (117), the *a priori* constraint [85]

$$b - a_x/2 = \lambda(t), \quad (318)$$

and the linearizing transformation of (G25) defines the link between τ and ψ

$$\frac{\tau_x}{\tau} = \frac{\lambda(t)}{Y_{1,x} + (1/2)Y_1^2 - a/2}, Y_1 = \frac{\psi_x}{\psi}. \quad (319)$$

This generates fourteen determining equations (150) (i.e. the same order of magnitude as for the (G5) assumption) in the basis $(\psi_x/\psi, \psi_{xx}/\psi - ((\psi_x/\psi)^2 + b_x\psi_x/(b\psi) - a)/2)$; they are solved as follows (g_k denotes an arbitrary integration function, λ an arbitrary integration constant) [90]

$$\begin{aligned} E_{4,4} &: c = 9\lambda(t)/\beta, \\ E_{3,5} &: a = -6V_x/\alpha, \\ E_{1,5} &: d = (3V_{xxx}/\alpha - (6V_x/\alpha)^2)/\beta, \\ E_{0,5} &: \lambda(t) = \lambda \text{ independent of } t, \\ E_{0,6} &: \text{pKK}(V) = 0, \\ E_{1,4} &: e = g_2(t) + (36\lambda V_x/\alpha + 72V_x V_{xx}/\alpha^2 + 3V_{xxx}/\alpha)/\beta, \\ X_1 &: b = g_4(t) - 3V_{xx}/\alpha + V_{xxx}/(3\alpha), \\ X_0 &: g_4 = \lambda, \end{aligned} \quad (320)$$

and the result is, with $U = V_x$,

$$b = \lambda - 3U_x/\alpha, \quad (321)$$

$$a = -6U/\alpha, \quad (322)$$

$$\partial_x \text{Log } \tau = \frac{\lambda}{\psi_{xx}/\psi - (1/2)(\psi_x/\psi)^2 + 3(U/\alpha)}. \quad (323)$$

$$L_1 = \partial_x^3 + 6\frac{U}{\alpha}\partial_x + 3\frac{U_x}{\alpha} - \lambda, \quad (324)$$

$$L_2 = \beta\partial_t - 9\lambda\partial_x^2 + \left(3\frac{U_{xx}}{\alpha} + 36\frac{U^2}{\alpha^2}\right)\partial_x - 3\frac{U_{xxx}}{\alpha} - 72\frac{UU_x}{\alpha^2} - 36\lambda\frac{U}{\alpha}, \quad (325)$$

$$\beta[L_1, L_2] = (6/\alpha)\text{KK}(U)\partial_x + (3/\alpha)\text{KK}(U)_x. \quad (326)$$

This is the Lax pair given by Kaup [65]. The integration of the first-order ODE (323) *modulo* the Lax pair yields the DT given by Levi and Ragnisco [74] :

$$\tau = \psi\psi_{xx} - (1/2)\psi_x^2 + 3(U/\alpha)\psi^2, \quad \tau_x = \lambda\psi^2. \quad (327)$$

Although the relation $\tau_x/\psi^2 = \text{constant}$ is the same as in the case of KdV (see Eq. (4.14) in Ref. [119]), it cannot be taken as an *a priori* assumption, it is the result of the method.

Starting from the *vacuum* solution $U = 0$, the general solution ψ of $L_1\psi = 0, L_2\psi = 0$,

$$\psi = c_1 e^{Kx+9K^5t/\beta} + c_2 e^{jKx+9j^2K^5t/\beta} + c_3 e^{j^2Kx+9jK^5t/\beta}, \quad (328)$$

$$j^3 = 1, \quad K^3 = \lambda,$$

in which c_1, c_2, c_3, K are arbitrary complex constants, leads by (141) to the one-soliton solution of Kaup [65]

$$u = (\alpha/2)\partial_x^2 \text{Log}(2 + \cosh(k/2)(x - (k/2)^4t/\beta)), \quad k \in \mathcal{R} \quad (329)$$

for the choice $(c_1, c_2, c_3) = (0, j^2, -j), K^2 = -k^2/12$, which corresponds to the entire function

$$\tau = -(k^2/12)(2 + \cosh(k/2)(x - (k/2)^4t/\beta))e^{(k/2)(x+(k/2)^4t/\beta)}. \quad (330)$$

Let us now obtain the auto-BT of KK, by an elimination. In order to perform this elimination easily, it is convenient to choose one of the two components of the pseudopotential \mathbf{Y} so as to characterize the DT,

$$\text{KK} : \frac{2(v - V)}{\alpha} = \frac{\tau_x}{\tau} = Z. \quad (331)$$

The chosen equivalent system is the system satisfied by (Y_1, Z)

$$Y_1 = \frac{\psi_x}{\psi}, \quad Z = \frac{\tau_x}{\tau}, \quad (332)$$

$$Y_{1,x} = -Y_1^2/2 + \lambda Z^{-1} - 3U/\alpha, \quad (333)$$

$$Z_x = 2Y_1 Z - Z^2, \quad (334)$$

$$\begin{aligned} \beta Y_{1,t} = & [9\lambda Y_1^2/2 - (3U_{xx}/\alpha + 36(U/\alpha)^2)Y_1 + 9\lambda^2 Z^{-1} \\ & + 3U_{xxx}/\alpha + 72UU_x/\alpha^2 + 9\lambda U/\alpha]_x, \end{aligned} \quad (335)$$

$$\begin{aligned} \beta Z_t = & [18\lambda U/\alpha + 9\lambda^2 Z^{-1} + 9\lambda Y_1^2 \\ & + (45(U/\alpha)^2 + 6(U_{xx}/\alpha) - 18(U_x/\alpha)Y_1 \\ & + 27(U/\alpha)Y_1^2 + (9/4)Y_1^4)Z]_x. \end{aligned} \quad (336)$$

The BT then arises from the elimination of Y_1 between (333), (334) and (336) (Eq. (335) must be discarded), which results in the two equations for $Z = Y$,

$$Y_{xx} - (3/4)Y_x^2/Y + 3YY_x/2 + Y^3/4 + 6(U/\alpha)Y - 2\lambda = 0, \quad (337)$$

$$\begin{aligned} \beta Y_t - (3/16)[3Y^5 + 15YY_x^2 + 30Y^2Y_{xx} + 8Y_{xxxx} \\ + 30(Y^3 + 2Y_{xx})(Y_x + 4V_x/\alpha) + 60Y(Y_x + 4V_x/\alpha)^2 \\ + 30Y_x(Y_{xx} + 4V_{xx}/\alpha) + 20Y(Y_{xxx} + 4V_{xxx}/\alpha)]_x = 0, \end{aligned} \quad (338)$$

$$\beta((Y_{xx})_t - (Y_t)_{xx})/Y = -(6/\alpha)KK(U), \quad (339)$$

followed by the substitution $Y = 2(v - V)/\alpha$, [90]

$$\begin{aligned} (v - V)_{xx}/\alpha - (3/4)(v - V)_x^2/(\alpha(v - V)) \\ + 3(v - V)(v + V)_x/\alpha^2 + (v - V)^3/\alpha^3 - \lambda = 0, \end{aligned} \quad (340)$$

$$\begin{aligned} \beta(v - V)_t/\alpha - (3/2)[2(v - V)_{xxxx}/\alpha + 60(v - V)^3(v + V)_x/\alpha^4 \\ + 12(v - V)^5/\alpha^5 + (10(v - V)(v + V)_{xxx} + 30(v + V)_x(v - V)_{xx} \\ + 15(v - V)_x(v + V)_{xx})/\alpha^2 + (30(v - V)^2(v - V)_{xx} \\ + 60(v - V)(v + V)_x^2 + 15(v - V)(v - V)_x^2/\alpha^3]_x = 0. \end{aligned} \quad (341)$$

The simple form of the conservative equations (312) and (341) results from the addition of suitable differential consequences of (311) and (340). The x -part of the BT has already been given by Rogers and Carillo [103] for $\lambda = 0$.

If we write (333)–(334) in the variables $(Y_1, Z_2 = Z^{-1})$,

$$Y_{1,x} = -Y_1^2/2 + \lambda Z_2 - 3U/\alpha, \quad (342)$$

$$Z_{2,x} = -2Y_1 Z_2 + 1, \quad (343)$$

both systems (124)–(125) and (342)–(343) are coupled Riccati systems, with the difference that the transformation from Y_1 to ψ_x/ψ is a point transformation while the one from Z_2 to ψ_x/ψ is a contact one. Thus, the Riccati system (124)–(125) is in the classification of linearizable coupled Riccati systems

given by Lie (this is the projective one), while the Riccati system (342)–(343) is outside it.

A minor open problem is to find a bilinear BT for SK equation (the one given in Ref. [90] contains a nonbilinear term).

7.3 Nonintegrable equations, second scattering order

Strictly speaking, nonintegrable equations have no associated scattering order. What is meant in the title of this section is that one assumes a given scattering order to process some nonintegrable PDEs.

For algebraic PDEs in two variables, particular solutions in which (S, C) are constant are quite easy to find. They correspond to solutions u polynomial in $\tanh \frac{k}{2}(x - ct - x_0)$. The privilege of \tanh is to be the general solution of the *unique* first order first degree nonlinear ODE with the PP, namely the Riccati equation $\tanh' + \tanh^2 + S/2$, in the particular case $S = \text{constant}$.

A characteristic feature of nonintegrable equations is the absence of a BT. Therefore, the iteration of Section 2 can only generate a finite number of new solutions [29,9], this will be seen on examples.

The Kuramoto-Sivashinsky equation

It is worth to handle this example in detail because it exhibits all the features of what should be done and more importantly of what should *not* be done when solving truncation equations.

The equation of Kuramoto and Sivashinsky (3) (notation $\mu = 19\mu'$) possesses a single family [94,49]

$$p = -3, \quad q = -7, \quad u_0 = 120\nu, \quad \text{indices } -1, 6, \frac{13 \pm i\sqrt{71}}{2}, \quad \mathcal{D} = 60\nu\partial_x^3 + 60\mu'\partial_x. \quad (344)$$

and the orthogonality condition at index 6 is satisfied. Since equation (3) is a conservation law, we therefore study it on its potential form

$$E \equiv v_t + \frac{v_x^2}{2} + \mu v_{xx} + \nu v_{xxxx} + G(t) = 0, \quad u = v_x, \quad (345)$$

which has the unique family

$$p = -2, \quad q = -6, \quad v_0 = -60\nu, \quad \text{indices } -1, 2, \frac{13 \pm i\sqrt{71}}{2}, \quad \mathcal{D} = 60\nu\partial_x^2 + 60\mu'. \quad (346)$$

Although the no-log condition at index $i = 2$ is not satisfied, the ψ -series

$$v = -60\nu\chi^{-2} + 60\mu' \text{Log } \psi + v_2 + 0(\chi), \quad v_2 \text{ arbitrary function}, \quad (347)$$

in which the gradients of ψ and χ are given by (68)–(69) and (64)–(65), contains one logarithm only, which cancels by derivation.

The one-family truncation assumption is

$$v_T = v_0\chi^{-2} + v_1\chi^{-1} + v_{02}\text{Log}\psi + v_2, \quad v_{02} \text{ constant} \quad (348)$$

equivalent to a truncated series $(-3 : 0)$ for u . Substituting (348) in (345) and eliminating any derivative of χ and ψ , one obtains

$$E = \sum_{j=0}^6 E_j \chi^{j-6}. \quad (349)$$

Together with the identity (62), this defines a system of eight equations in the six unknowns $(v_0, v_1, v_{02}, v_2, S, C)$.

Equations $j = 0, 1, 2$ are solved for v_0, v_1, v_{02}, v_2 exactly as in the Painlevé test and yield the values in (347). The next five equations ($j = 3, 4, 5, 6$ and (62)) now read [29]

$$E_3 \equiv 120\nu(-C + 15\nu S_x + v_{2,x}) = 0, \quad (350)$$

$$E_4 \equiv 60(-6\nu^2 S_{xx} - 4\nu^2 S^2 - 20\mu'\nu S + 2\nu C_x + 11\mu'^2) = 0, \quad (351)$$

$$E_5 \equiv \frac{S}{2}E_3 + 60(-\mu'C + 20\mu'\nu S_x - 2\nu^2 S S_x + \nu^2 S_{xxx} - \nu C_{xx} + \mu'v_{2,x}) = 0, \quad (352)$$

$$E_6 \equiv E(v_2) + 30(\mu'C_x - 19\mu'^2 S - \mu'\nu(20S^2 + S_{xx}) - \nu^2(4S^3 + 3S_x^2 + 4SS_{xx})) = 0,$$

$$X \equiv S_t + C_{xxx} + 2C_x S + C S_x = 0. \quad (353)$$

The principles to be obeyed during the resolution are the following.

1. Never increase the differential order of a given variable. On the contrary, solve for the higher derivatives in terms of the lower ones, and substitute the result, as well as its differential consequences, in the remaining equations.
2. Never integrate a differential equation, unless it is just a total derivative. On the contrary, perform an *algebraic* resolution.
3. Never solve for a function of, say, one variable as an expression in several variables.
4. Close the solution by exhausting all Schwarz cross-derivative conditions.

This computation is systematic, and its algorithmic version is known as the construction of a differential Groebner basis [79,6].

The full system is split into (E_3, E_4, E_5) , independent of ∂_t , and (E_6, X) , explicitly depending on ∂_t . The subsystem (E_3, E_4) is first solved, according to rule 1, as a Cramer system for $(v_{2,x}, S_{xx})$. After substitution of $(v_{2,x}, S_{xx})$ and their derivatives in all the other equations, equation E_5 is solved, according to rule 1, for C_{xx} and the result is recognized as being an x -derivative. This allows us to solve for C_x after the introduction of an arbitrary integration function λ of t only. As to (E_6, X) , they are solved, according to rule 1,

as a Cramer system in variables involving only t -derivatives, namely $(v_{2,t}, S_t)$, for expressions independent of ∂_t . To summarize this first stage, the original system is now equivalent to

$$S_{xx} = -\frac{3}{2}S^2 - \frac{5\mu'}{2\nu}S + \frac{\mu'^2}{8\nu^2}(\lambda + 22), \quad (354)$$

$$C_x = -\frac{5\nu}{2}S^2 + \frac{5\mu'}{2}S + \frac{\mu'^2}{8\nu}(3\lambda + 22), \quad (355)$$

$$v_{2,x} = C - 15\nu S_x, \quad (356)$$

$$\begin{aligned} \frac{v_{2,t}}{\nu} = & -\frac{1}{\nu}G(t) + \frac{1243\mu'^3}{2\nu^2} + 16\frac{\mu'^3}{\nu^2}\lambda + 10\frac{\mu'^2}{\nu}(\lambda - 2)S \\ & + 110\mu'S^2 - \frac{1}{2\nu}C^2 + 15CS_x - \frac{125\nu}{2}S_x^2, \end{aligned} \quad (357)$$

$$\begin{aligned} S_t = & -\frac{5\mu'^3}{16\nu^2}(\lambda + 22) - \frac{\mu'^2}{8\nu}(\lambda - 116)S - \frac{55\mu'}{4}S^2 - \frac{5\nu}{2}S^3 \\ & - CS_x + 5\nu S_x^2. \end{aligned} \quad (358)$$

One equation, and only one, namely (354), is an ODE. Integrating it as an elliptic ODE for S [29] would create useless subcases and complications and should, according to rule 2, *not* be done. This ODE should also *not* be replaced by its first integral, because the integrating factor S_x could be, and will indeed be, zero. According to rule 3, it is also forbidden to eliminate $\lambda(t)$ by solving e.g. (355) for it. The only thing to do is (rule 4) to close this solution by cross-differentiation. There are two such conditions:

$$\begin{aligned} (S_{xx})_t - (S_t)_{xx} \equiv & -\frac{\mu'^4}{64\nu^3}(3\lambda^2 + 308\lambda + 5324) + \frac{5\mu'^3}{8\nu^2}(15\lambda + 374)S \\ & + \frac{25\mu'^2}{8\nu}(\lambda - 16)S^2 - \frac{165\mu'}{2}S^3 - \frac{75\nu}{4}S^4 \\ & + \frac{\mu'^2}{8\nu^2}\lambda' + \frac{85\mu'}{2}S_x^2 + 25\nu SS_x^2 = 0, \end{aligned} \quad (359)$$

$$\begin{aligned} (v_{2,t})_x - (v_{2,x})_t \equiv & -C_t - \frac{\mu'^2}{8\nu}(3\lambda + 22)C - \frac{5\mu'}{2}SC + \frac{5\nu}{2}S^2C \\ & + \frac{15\mu'^2}{4}(3\lambda + 71)S_x - 255\mu'\nu SS_x \\ & - 150\nu^2 S^2 S_x = 0. \end{aligned} \quad (360)$$

The latter is solved for C_t and provides a third cross-derivative condition

$$\begin{aligned} \frac{(C_x)_t - (C_t)_x}{\nu} \equiv & -\frac{\mu'^4}{64\nu^3}(81\lambda^2 + 4028\lambda + 47476) \\ & + \frac{5\mu'^3}{8\nu^2}(101\lambda + 2322)S + \frac{5\mu'^2}{8\nu}(55\lambda + 96)S^2 \\ & - \frac{1415\mu'}{2}S^3 - \frac{825\nu}{4}S^4 \end{aligned}$$

$$+\frac{3\mu'^2}{8\nu^2}\lambda' + \frac{535\mu'}{2}S_x^2 + 275\nu SS_x^2 = 0. \quad (361)$$

This ends the linear part of the resolution, and now comes the nonlinear part (algebraic Groebner). The two remaining equations (359) and (361), considered as nonlinear in the two unknowns (S_x, S) , imply without computation $S_x = 0$, which then allows one to solve (354) for the monomial S^2 as a polynomial in S of a smaller degree. Equations (359) and (361) thus become linear in S and λ' , and their resultant in λ' factorizes as a product of *linear* factors

$$(\lambda + 33)(\lambda - 11 + \frac{40\nu}{\mu'}S) = 0. \quad (362)$$

The first factor yields no solution. The second one provides the unique solution

$$\left(S - \frac{\mu}{38\nu}\right)\left(S + \frac{11\mu}{38\nu}\right) = 0, \quad C = \text{arbitrary constant } c \quad (363)$$

and it leads to the two-parameter (c, x_0) solution (4). The two equations (363) represent the SME.

If one performs the iteration of Section 2, starting from $u = c$, one generates the solitary wave (4) and no more [29].

The reduction $u(x, t) = c + U(\xi)$, $\xi = x - ct$ of the PDE (3) yields the ODE

$$\nu U''' + \mu U' + U^2/2 + K = 0, \quad K \text{ arbitrary}, \quad (364)$$

with the indices $-1, (13 \pm i\sqrt{71})/2$. Due to the two irrational indices, the *general analytic solution* (see definition in Section 9.3) can only depend on one arbitrary constant. This one-parameter solution, whose local expansion contains no logarithm, is known globally only for $K = -450\nu k^2/(19^2\mu)$, this is (4), but its closed form expression for any K is still an open problem.

Being autonomous, the ODE (364) is equivalent to the nonautonomous second order ODE for $V(U)$

$$V = \frac{dU}{d\xi} : \nu \frac{d^2(V^2)}{dU^2} + 2\mu + \frac{U^2 + 2K}{V} = 0, \quad (365)$$

an equation which has been studied from the Hamiltonian point of view [7].

7.4 Nonintegrable equations, third scattering order

An example is given in the Section 9.3.

8 Two common errors in the one-family truncation

Two errors are frequently made in the method of section 7.

8.1 The constant level term does not define a BT

Consider the one-family truncation as done by WTC (the subscript T means “truncated”)

$$u_T^{\text{WTC}} = \sum_{j=0}^{-p} u_j^{\text{WTC}} \varphi^{j+p} \tag{366}$$

in which φ is the function defining the singularity manifold.

In the WTC truncation, one considers three solutions of the PDE

1. the lhs u_T^{WTC} of the truncation (366),
2. the “constant level” coefficient u_{-p}^{WTC} ,
3. the field U which appears in the Lax pair after the successful completion of the method.

The frequently encountered argument “The constant level coefficient u_{-p}^{WTC} also satisfies the PDE, therefore one has obtained a BT” is wrong. This is obvious, since nonintegrable PDEs, which have no BT, nevertheless have this property. One can check it by taking the explicit example of a nonintegrable PDE [29].

A hint that the above argument might be wrong is the fact, observed on all successful truncations, that the U in the Lax pair is *never* u_{-p}^{WTC} . Let us prove this fact, with the homographically invariant analysis [19]. The truncation of the same variable in the invariant formalism is

$$u_T = \sum_{j=0}^{-p} u_j \chi^{j+p}, \tag{367}$$

in which χ is given by (58). This u_T depends on the movable constant φ_0 and one has

$$\begin{cases} u_T^{\text{WTC}} = u_T(\varphi_0 = 0) \\ u_{-p}^{\text{WTC}} = u_T(\varphi_0 = \infty). \end{cases} \tag{368}$$

Since the results of the truncation do not depend on the movable constant φ_0 , this proves that the lhs u_T^{WTC} of the truncation and the constant level coefficient u_{-p}^{WTC} are not considered as distinct by the singular manifold method. Since the U in the Lax pair cannot be the truncated u (otherwise one would not have a Darboux transformation), this ends the proof.

8.2 The WTC truncation is suitable iff the Lax order is two

We mean the truncation as originally introduced, not its updated version of Section 7.

When the Lax pair has second order, everything is consistent. When the Lax pair has a higher order, e.g. three, the original method, as well as its

original invariant version [86], presents the following inconsistency. In a first stage, it generates the $-q + p$ equations $E_j(S, C, U) = 0$ of formula (147), which intrinsically correspond to a *second*-order scattering problem (and this is precisely the inconsistency), and in a second stage it injects in each of these $-q + p$ equations a link between (S, C) and the scalar field ψ of the Lax pair of *higher* order, thus generating determining equations which are hybrid between the second order and the higher one. The first nearly correct treatment has been made in Ref. [87].

For the same reason, in order to obtain the Lax pair when its order is higher than two, it is also inconsistent to consider the so-called *singular manifold equation* (SME) [119,19,101], defined in Section 7.1. When the Lax order is three, the correct extension of this SME notion would be the set of three relations on (a, b, c, d) resulting from the elimination of U between the four coefficients of the Lax pair (e is derivable from (122) so we discard it), but this seems of little interest.

Although these inconsistencies may still provide the full result for some “robust” equations (Boussinesq [117], Sawada-Kotera [115], Hirota-Satsuma [86]), there do exist equations for which it leads to a failure, and the Kaup-Kupershmidt equation [90] is one of them.

9 The singular manifold method applied to two-family PDEs

By two-family, we mean two opposite families.

This includes also the one-family truncation as a particular case.

When the base member of the hierarchy of integrable equations has more than a single family, these families usually come by pairs of opposite singular part operators, just like (P2)–(P6). Examples are enumerated at the end of Section 5.2. Then the sum of the two opposite singular parts

$$\mathcal{D} \text{Log } \tau_1 - \mathcal{D} \text{Log } \tau_2 \tag{369}$$

only depends on the variable

$$Y = \frac{\tau_1}{\tau_2}. \tag{370}$$

The current status of the method [88,101], which used to be called the two-singular manifold method [88], is as follows. Most of the method for one-family equations still applies, with the difference that it is much more convenient to represent the Lax pairs in a Riccati form than in a scalar linear form. Let us restrict here to second-order scattering problems (for the third order case, see Section 9.2) and to identity links (146) between the two τ and the two ψ functions. Then Y satisfies a Riccati system and, as explained in Section 6.1, its most general expression is given by (66).

In the first step, τ is simply replaced by Y in the assumption (141) for a DT.

In the second step, the scattering problem is represented by the Riccati system satisfied by Y , whose coefficients depend on (S, C, A, B) .

The fourth step contains the main difference. Rather than truncating u at the level $j = -p$, one truncates it at the level $j = -2p$ [88,101], in order to implement the two movable singularities $\tau_1 = 0$ and $\tau_2 = 0$. So the truncation is [101] (for second order Lax pairs only)

$$u = \mathcal{D} \operatorname{Log} Y + U, \quad (371)$$

$$Y^{-1} = B(\chi^{-1} + A), \quad (372)$$

$$E(u) = \sum_{j=0}^{-2q} E_j(S, C, A, B, U) Y^{j+q}, \quad (373)$$

$$\forall j \ E_j(S, C, A, B, U) = 0, \quad (374)$$

in which nothing is imposed on U .

Let us remark that the relation $A \neq 0$ does not characterize two-family PDEs, see the Liouville case in Section 9.1.

9.1 Integrable equations with a second order Lax pair

The sine-Gordon equation

The sine-Gordon equation is defined for convenience as the case $a_1 \neq 0, a_2 = 0$ of the equation (168). Although not algebraic in u , it becomes algebraic in e^u and it possesses two opposite families (opposite in the field u), both with $p = -2, q = -2$

$$e^u \sim -(2/\alpha)\varphi_x\varphi_t(\varphi - \varphi_0)^{-2}, \text{ indices } (-1, 2), \mathcal{D} = (2/\alpha)\partial_x\partial_t. \quad (375)$$

$$e^{-u} \sim (2/a_1)\varphi_x\varphi_t(\varphi - \varphi_0)^{-2}, \text{ indices } (-1, 2), \mathcal{D} = -(2/a_1)\partial_x\partial_t. \quad (376)$$

The resulting DT assumption

$$e^u + (a_1/\alpha)e^{-u} = (2/\alpha)\partial_x\partial_t \operatorname{Log} Y + \tilde{W}, \ E(u) = 0 \quad (377)$$

with Y defined by (370), can be integrated twice due to the special form of the PDE, resulting in

$$u = -2 \operatorname{Log} Y + W, \ E(u) = 0, \quad (378)$$

in which nothing is imposed on W (we use W to reserve the symbol U for future use). For $a_1 = 0$, this truncation is what was called in Section 7.1 the second truncation of Liouville equation.

The five determining equations in the unknowns (S, C, A, B, W) are [101,37]

$$E_0 \equiv \alpha B^2 e^W - 2C = 0, \quad (379)$$

$$E_1 \equiv 2(C_x + 2AC) = 0, \quad (380)$$

$$E_2 \equiv 0, \text{ (Fuchs index)} \quad (381)$$

$$E_3 \equiv -\sigma_t - \sigma(C_x + 2AC) = 0, \quad (382)$$

$$E_4 \equiv \sigma(C\sigma + (C_x + 2AC)_x)/2 + a_1 B^{-2} e^{-W} = 0, \quad (383)$$

with the abbreviation

$$\sigma = S + 2A^2 - 2A_x, \quad (384)$$

and, together with the cross-derivative condition (62), they are solved as usual by ascending values of j

$$E_0 : B^2 e^W = \frac{2}{\alpha} C, \quad (385)$$

$$E_1 : A = -\frac{1}{2} (\text{Log } C)_x, \quad (386)$$

$$E_3 : S = -F(x) + \frac{C_x^2}{2C^2} - \frac{C_{xx}}{C}, \quad (387)$$

$$E_4 : CC_{xt} - C_x C_t + F(x)C^3 + a_1 \alpha F(x)^{-1} C = 0, \quad (388)$$

$$X : a_1 F'(x) = 0. \quad (389)$$

in which F is a function of integration. For sine-Gordon, $F(x)$ must be a constant

$$F(x) = 2\lambda^2. \quad (390)$$

In the Liouville case, for which the truncation imposes no restriction on $F(x)$, let us also require that $F(x)$ be a constant. Then, for both equations, $\text{Log } C$ is proportional to a second solution U of the PDE

$$C = \frac{\alpha}{2} \lambda^{-2} e^U, \quad E(U) = 0, \quad (391)$$

and one has obtained the Darboux transformation

$$u = -2 \text{Log } y + U, \quad y = \lambda B Y, \quad (392)$$

in which y satisfies the Riccati system

$$y_x = \lambda + U_x y - \lambda y^2, \quad (393)$$

$$y_t = -\frac{\alpha}{2} \lambda^{-1} (e^U + (a_1/\alpha) e^{-U} y^2), \quad (394)$$

$$(\text{Log } y)_{xt} - (\text{Log } y)_{tx} = E(U). \quad (395)$$

The linearization

$$y = \psi_1/\psi_2 \quad (396)$$

yields the second-order matrix Lax pair

$$(\partial_x - L) \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = 0, \quad L = \begin{pmatrix} U_x/2 & \lambda \\ \lambda & -U_x/2 \end{pmatrix}, \quad (397)$$

$$(\partial_t - M) \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = 0, \quad M = -(\alpha/2)\lambda^{-1} \begin{pmatrix} 0 & e^U \\ -(a_1/\alpha)e^{-U} & 0 \end{pmatrix}. \quad (398)$$

The auto-BT (classical for sine-Gordon, Ref. [81] for Liouville) results from the substitution $y = e^{-(u-U)/2}$ into (393)–(394)

$$(u + \tilde{U})_x = -4\lambda \sinh \frac{u - \tilde{U}}{2}, \quad (399)$$

$$(u - \tilde{U})_t = \lambda^{-1} \left(\alpha e^{(u+\tilde{U})/2} + a_1 e^{-(u+\tilde{U})/2} \right). \quad (400)$$

It coincides in the sine-Gordon case with the one given earlier, equations (9)–(10). The ODE part (393) of the BT is a Riccati equation.

The SME is [101]

$$S + C^{-1}C_{xx} - \frac{1}{2}C^{-2}C_x^2 + 2\lambda^2 = 0, \quad (401)$$

and it coincides, but this is not generic, with the one [116,19] obtained from the (incorrect) truncation in χ .

Remarks.

1. The reason for the presence of the apparently useless parameter B in the definition (66) is to allow the precise correspondence (146)

$$\tau_1 = \psi_1, \quad \tau_2 = \psi_2 \quad (402)$$

for some choice of B , namely

$$B = \lambda^{-1}, \quad y = Y, \quad W = U. \quad (403)$$

2. In the Liouville case $a_1 = 0$, this is an example of a PDE with only one family and a nonzero value of A .

The modified Korteweg-de Vries equation

This PDE has the same scattering problem as sine-Gordon, so the computation should be, and indeed is, quite similar to that for sine-Gordon.

Since this PDE has the conservative form

$$\text{mKdV}(w) \equiv bw_t + (w_{xx} - 2(w - \beta)^3/\alpha^2 + 6vw)_x = 0, \quad w = r_x, \quad (404)$$

it is technically cheaper to process its potential form

$$\text{p-mKdV}(r) \equiv br_t + r_{xxx} - 2(r_x - \beta)^3/\alpha^2 + 6\nu(r_x - \beta) + F(t) = 0. \quad (405)$$

Its invariance under the involution $w - \beta \mapsto -(w - \beta)$ provides an elegant way [76] to derive the BT of the KdV equation and its hierarchy. Although the constants β and ν could be set to zero by a transformation on (r, x, t) preserving the PP, it is convenient to keep them nonzero, for reasons explained at the end of this Section. This last PDE admits the two opposite families (α is any square root of α^2)

$$p = 0^-, \quad q = -3, \quad r \sim \alpha \text{Log } \psi, \quad \text{indices } (-1, 0, 4), \quad \mathcal{D} = \alpha. \quad (406)$$

The truncation is defined by

$$r = \alpha \text{Log } Y + R, \quad (407)$$

with (371)–(374), and this generates five equations $E_j = 0$ [101], with the notation (384) for σ

$$E_1 \equiv 6\alpha A - 6((R - \alpha \text{Log } B)_x - \beta) = 0, \quad (408)$$

$$E_2 \equiv \alpha(2A_x + 4A^2 - bC - 2\sigma + 6\nu) - \alpha^{-1}(E_1 + 6\alpha A)^2/6 = 0, \quad (409)$$

$$E_3 \equiv \text{p-mKdV}(R - \alpha \text{Log } B) - (3/2)\alpha^{-1}\sigma_x + (\sigma - 4A^2 - (1/3)\alpha^{-1}E_{1,x} - 2A_x)E_1 - 2AE_{1,x} - 2AE_2 - E_{2,x} \quad (410)$$

$$E_4 \equiv \text{expression vanishing with } E_1, E_2, E_3, E_5, \quad (411)$$

$$E_5 \equiv (3/4)\alpha\sigma\sigma_x + (1/4)\sigma^2 E_1 = 0, \quad (412)$$

$$X \equiv S_t + C_{xxx} + 2C_x S + C S_x = 0. \quad (413)$$

They depend on (R, B) only through the combination $R - \alpha \text{Log } B$. Equation $j = 4$ is a differential consequence of equations $j = 1, 2, 3, 5$, because 4 is a Fuchs index, and the other equations have been written so as to display how they are solved:

$$E_1 : A = \alpha^{-1}((R - \alpha \text{Log } B)_x - \beta), \quad (414)$$

$$E_5 : \sigma = -2(\lambda(t)^2 - \nu), \quad \lambda \text{ arbitrary function}, \quad (415)$$

$$E_2 : bC = 2A_x - 2A^2 + 4\lambda(t)^2 + 2\nu, \quad (416)$$

$$E_3 : \text{p-mKdV}(R - \alpha \text{Log } B) = 0, \quad (417)$$

$$X : \lambda'(t) = 0. \quad (418)$$

Thus, their general solution can be expressed in terms of a second solution W of the mKdV equation (404) and an arbitrary complex constant λ [101]

$$\begin{aligned} W &= (R - \alpha \text{Log } B)_x, \quad A = (W - \beta)/\alpha, \\ bC &= 2W_x/\alpha - 2(W - \beta)^2/\alpha^2 + 2\nu + 4\lambda^2, \\ S &= 2W_x/\alpha - 2(W - \beta)^2/\alpha^2 + 2\nu - 2\lambda^2, \end{aligned} \quad (419)$$

and the cross-derivative condition $X_1 = 0$ (Eq. (106)), equivalent to the mKdV equation (404) for W , proves that one has obtained a Darboux transformation and a Lax pair.

The SME, obtained by the elimination of W between S and C ,

$$bC - S - 6\lambda^2 = 0, \tag{420}$$

is identical to that of the KdV equation (216).

The auto-BT of mKdV is obtained by the substitution

$$\text{Log}(BY) = \alpha^{-1} \int (w - W) dx \tag{421}$$

in the two equations for the gradient of $y = BY$

$$\frac{y_x}{y} = \lambda \left(\frac{1}{y} - y \right) - 2 \frac{W - \beta}{\alpha}, \tag{422}$$

$$b \frac{y_t}{y} = \frac{1}{y} \left(-4\lambda \frac{W - \beta}{\alpha} + \left(2 \frac{(W - \beta)^2}{\alpha^2} + 2 \frac{W_x}{\alpha} - 4\lambda^2 \right) y \right)_x. \tag{423}$$

In the same manner as in the KdV truncation, these two Riccati equations can also be interpreted as the hetero-BT between the mKdV equation and the PDE satisfied by the pseudopotential y , called the Chen-Calogero-Degasperis-Fokas PDE.

The nonlinear Schrödinger equation

For the AKNS system of two second order equations in (u, v) (whose reduction $\bar{u} = v$ is NLS, see (39)), no two-family truncation has yet been defined which strictly follows the method and provides the desired result. It should be noted that the fourth order equation for u resulting from the elimination of v , known as the Broer-Kaup equation or classical Boussinesq system, admits a two-family truncation without any problem [38].

The full result (DT, BT) can be found [34] for the AKNS system by performing the one-family truncation [117] and then applying four involutions to the result of Weiss.

A second open problem for this PDE is that its bilinear BT is not yet known.

9.2 Integrable equations with a third order Lax pair

In principle, there is no additional difficulty to extend the method to a scattering order higher than two. A good equation to process would be the modified Boussinesq equation

$$E \equiv \begin{cases} -u_t + (v - (3/2)a^2u^2)_x = 0, \\ -v_t - 3a^2(u_{xx} - uv + a^2u^3)_x = 0, \end{cases} \tag{424}$$

which has two opposite families

$$u \sim (2/a)\chi^{-1}, \quad v \sim 6\chi^{-2} \quad (425)$$

and a third order Lax pair like the Boussinesq equation.

The one-family assumption [47]

$$u = U + (2/a)\partial_x \text{Log } \tau, \quad v = V - 6\partial_x^2 \text{Log } \tau, \quad E(u, v) = E(U, V) = 0, \quad (426)$$

with the identity link $\tau = \psi$ and the choice of the scalar Lax pair (117)–(118), already leads to the solution

$$\begin{aligned} f &= -(3/2)aU, \quad a = (V - 3a^2U^2 - 3aU_x)/4, \quad b = \lambda, \\ c &= -3a, \quad d = -3a^2U, \quad e = 0. \end{aligned} \quad (427)$$

Despite this success, it would be more consistent with the two-family structure to process this PDE with a two-family assumption, removing in passing the restriction $E(U, V) = 0$ in (426). This could make the coefficients (f, a, b) linear in (U, V) , which is not the case in (427).

Table 1 summarizes, for a sample of PDEs, the currently best method to obtain its Lax pair, Darboux and Bäcklund transformations from a truncation.

9.3 Nonintegrable equations, second and third scattering order

A nonintegrable equation has no determined scattering order, so this section cannot be split according to the scattering order.

The KPP equation

The KPP equation (77) possesses the two opposite families (78)–(80) and it fails the test at index 4, so there can only exist particular solutions. Let us first review all the known solutions to this equation.

In addition to the notation (81), it is convenient to introduce the symmetric constant

$$a_1 = (2e_1 - e_2 - e_3)(2e_2 - e_3 - e_1)(2e_3 - e_1 - e_2)/(3d)^3 \quad (428)$$

and the entire function

$$\Psi_3 = \sum_{n=1}^3 C_n e^{k_n(x + (3/b)k_n t)}, \quad k_n = \frac{3e_n - s_1}{3d}, \quad C_n \text{ arbitrary}, \quad (429)$$

i.e. the general solution of the third order linear system (117)–(118) with constant coefficients [31]

$$(S) \equiv \begin{cases} \psi_{xxx} - 3a_2\psi_x - a_1\psi = 0, \\ b\psi_t - 3\psi_{xx} = 0. \end{cases} \quad (430)$$

Table 1. The relevant truncation for some 1 + 1-dimensional PDEs. The successive columns are : the usual name of the PDE (a p means the potential equation), its number of families (a * indicates that only one family is relevant, see details in Ref), the order of its Lax pair, the truncation variable(s), the link between τ and ψ , the singularity orders of u and $E(u)$, the Fuchs indices (without the ever present -1), the number of determining equations, the reference to the place where the right method was first applied (earlier references may be found in it). The “?” in the AKNS system entry (the one whose NLS is a reduction) means that the method has not yet been applied to it, see text.

Name	f	Lax	Trunc. var.	τ	$-p : -q$	indices	nb. det. eq.	Ref
Liouville	1		τ		0 : 2	2	3	[37]
KdV	1	2	χ	ψ	2 : 5	4, 6	2	[119]
AKNS eq.	1	2	χ	ψ	1 : 5	4, 6	3	[83]
p-mKdV	2	2	Y	ψ	0 : 3	0, 4	4	[101]
sine-Gordon	2	2	Y	ψ	0 : 2	2	4	[101]
Broer-Kaup	2	2	Y	ψ	0 : 4	0, 3, 4	4	[101]
pp-Boussinesq	1	3	$(\psi_x/\psi, \psi_{xx}/\psi)$	ψ	0 : 4	0, 1, 6	6	[87]
p-SK	1*	3	$(\psi_x/\psi, \psi_{xx}/\psi)$	ψ	1 : 6	1, 2, 3, 10	6	[90]
p-KK	1*	3	$(\psi_x/\psi, \psi_{xx}/\psi)$	$G_{25}(\psi)$	1 : 6	1, 3, 5, 7	14	[85]
Tzitzéica	1*	3	$(\psi_x/\psi, \psi_t/\psi)$	ψ	2 : 6	2	10	[37]
AKNS system	4	2	?		1 : 3, 1 : 3	0, 3, 4		[34]

Let us also denote (j, l, m) any permutation of $(1, 2, 3)$. Three distinct solutions are presently known.

The first solution is trigonometric, this is a *collision of two fronts* [66]

$$u = \frac{s_1}{3} + d\partial_x \text{Log} \Psi_3, \quad C_1 C_2 C_3 \neq 0 \quad (431)$$

which depends on two arbitrary constants $C_1/C_3, C_2/C_3$. For $C_j = 0, C_l C_m \neq 0$, it degenerates into three heteroclinic (i.e. with different limits at both infinities) *propagating fronts* which depend on one arbitrary constant x_0

$$u = \frac{e_l + e_m}{2} + d\frac{k}{2} \tanh \frac{k}{2}(x - ct - x_0), \quad (432)$$

$$k^2 = (k_l - k_m)^2, \quad c = -3(k_l + k_m)/b.$$

The second solution is elliptic [10],

$$u = s_1/3 + d\psi_x \sqrt{\wp(\psi)}, \quad \psi = \Psi_3, \quad g_3 = 0, \quad g_2 \text{ arbitrary}, \quad a_1 = 0, \quad (433)$$

it only exists under the constraint (codimension is one) that one root e_j be at the middle of the two others and it depends on the four arbitrary constants

C_1, C_2, C_3, g_2 . Its degeneracy $g_2 = 0$ (i.e. $\wp(\psi) = \psi^{-2}$) is the degeneracy $a_1 = 0$ of the collision of two fronts solution (431).

The third and last solution is the stationary elliptic solution $u(x)$

$$u(x) : -u'' + 2d^{-2}(u - e_1)(u - e_2)(u - e_3) = 0. \quad (434)$$

A trigonometric degeneracy bounded at infinity is made of the three homoclinic *stationary pulses*

$$u = e_j + \frac{e_l - e_m}{\sqrt{2}} \operatorname{sech} i \frac{e_l - e_m}{d\sqrt{2}}(x - x_0), \quad a_1 = 0, \quad 2e_j - e_l - e_m = 0, \quad (435)$$

it has codimension one and it depends on the arbitrary constant x_0 .

Let us now apply the various methods we have seen, in order to retrieve these solutions, namely

1. enforcement of one of the two no-log conditions (82),
2. enforcement of the two no-log conditions (82),
3. one-family truncation with a second order assumption,
4. one-family truncation with a third order assumption,
5. two-family truncation with a second order assumption,
6. two-family truncation with a third order assumption.

The single no-log condition (82) has two solutions. The first one $C = 0$ implies $u_t = 0$ and thus defines the reduction $u(x)$, i.e. the elliptic equation (434). The second one is a first order nonlinear PDE for $C(x, t)$, integrated by the method of characteristics as [9]

$$\begin{aligned} F(I_1, I_2) &= 0, \quad c_n = 3k_n/b = (3e_n - s_1)/(bd), \\ I_1 &= e^x(C - c_1)^{p_1}(C - c_2)^{p_2}(C - c_3)^{p_3}, \\ I_2 &= e^t(C - c_1)^{q_1}(C - c_2)^{q_2}(C - c_3)^{q_3}, \end{aligned} \quad (436)$$

in which p_n, q_n depend on e_j . Unless some specific choice of the arbitrary function F is made, or (x, t) are no more taken as the independent variables, one cannot integrate further the system (64)–(65) for χ and S .

The two no-log conditions (82) together, apart the already encountered solution $C = 0$, provide the two relations

$$\begin{aligned} a_1 &= 0, \quad e_1 = (e_2 + e_3)/2, \\ b^2 d^2 C^3 - (9/4)(e_2 - e_3)^2 C - 3bd^2(C_t + CC_x) &= 0, \end{aligned} \quad (437)$$

whose solution is similarly

$$\begin{aligned} a_1 &= 0, \quad F(I_1, I_2) = 0, \\ I_1 &= \frac{bC - (3(e_2 - e_3)/(2d))}{bC + (3(e_2 - e_3)/(2d))} e^{(e_2 - e_3)x/d}, \\ I_2 &= \frac{C^2}{(bC)^2 - (3(e_2 - e_3)/(2d))^2} e^{(e_2 - e_3)^2 t/(bd^2)}. \end{aligned} \quad (438)$$

The one-family truncation (142) with $\tau = \psi$ and the second order assumption (111)–(112) generates three determining equations $E_j(S, C, U) = 0, j = 1, 2, 3$. After solving the first one for U

$$U = s_1/3 - bdC/6, \quad (439)$$

the two remaining equations, with the ever present condition (62), are [21]

$$E_2 \equiv -6a_2 - S + (b^2/6)C^2 - bC_x = 0 \quad (440)$$

$$E_3 \equiv a_2bC - 2a_1 + bSC/2 - b^3C^3/108 \\ + S_x/2 - b^2C_t/6 + 2bC_{xx}/3 = 0. \quad (441)$$

The elimination of S and C_t yields a factorized equation

$$-b^2C_t + bC_{xx} - 2b^2CC_x + (4/9)(bC)^3 - 12a_2bC - 12a_1 = 0, \quad (442)$$

$$[bC_{xx} - b^2CC_x + b^3C^3/9 - 3a_2bC - 3a_1]C = 0. \quad (443)$$

The subcase $C = 0$, hence $S = -6a_2, a_1 = 0$, yields the degeneracy $a_1 = 0$ of the three fronts (432). In the other subcase, the system for C is linearizable into the third order system (S) (430) in which both ∂_x and ∂_t change sign, the generic solution (S, C) of (E_2, E_3) is therefore

$$bC = -3\partial_x \text{Log } \Psi_3(-x, -t), \quad S = -6a_2 + (bC)^2/6 - bC_x, \quad (444)$$

and there only remains to integrate (64)–(65) for χ or (60)–(61) for φ . Since the one-form $dx - Cdt$ possesses an integrating factor [9], the PDE (61) for φ can be integrated by the method of characteristics,

$$\varphi = \Phi(F), \quad F = \frac{\Psi_x + k_2\Psi}{\Psi_x + k_3\Psi} \frac{e^{-k_2x - k_2^2t/b}}{e^{-k_3x - k_3^2t/b}}. \quad (445)$$

Note that the cyclic permutation of the roots e_j is broken when going from (S, C) to φ . With the classical identity on Schwarzians

$$\{\varphi; x\} \equiv \{\Phi; F\}F_x^2 + \{F; x\}, \quad (446)$$

the third order ODE (60) for φ becomes

$$\{\Phi; F\} = 0, \quad (447)$$

which integrates as

$$\varphi = \Phi(F) = \frac{A_1F + A_2}{A_3F + A_4}, \quad A_j \text{ arbitrary constants, } A_1A_4 - A_2A_3 \neq 0. \quad (448)$$

The value of χ^{-1}

$$\chi^{-1} = \frac{F_x}{F - F_0} - \frac{F_{xx}}{2F}, \quad F_0 \text{ arbitrary constant,} \quad (449)$$

is again invariant under a cyclic permutation of the roots e_j , and the solution u finally obtained is (431).

The one-family truncation (142) with $\tau = \psi$ and the third order assumption (117)–(118) generates five determining equations (150), their straightforward resolution yields

$$u = d\partial_x \text{Log } \psi + s_1/3 + dU, \quad (450)$$

$$\begin{aligned} \psi_{xxx} + 3U\psi_{xx} - 3(a_2 - U^2 - U_x)\psi_x \\ - (a_1 + 3a_2U - U^3 - bU_t/2 - 6UU_x + U_{xx}/2)\psi = 0, \end{aligned} \quad (451)$$

$$b\psi_t - 3\psi_{xx} - 3U\psi_x - 6U^2\psi = 0, \quad (452)$$

in which U is constrained by two relations. But, since the coefficient f can be set to zero without loss of generality, the choice $U = 0$ represents the general solution (just like in [31] where constant values were assumed *ab initio* for the coefficients in (117)–(118)), and it represents again the collision of two fronts (431).

The contrast of difficulty between the second order assumption (laborious) and the third order assumption (immediate) is the signature that the good scattering order of KPP is three, despite the irrelevance of such a notion for nonintegrable equations.

The two-family truncation with a second order assumption (371)–(374) [30,31,100] generates five determining equations. Despite the factorized form of E_5 , we have not yet found their general solution. The three particular solutions for which (S, C, A) are constant provide immediately the three pulses (435), for they belong to the class of polynomials in \tanh and sech , generated by negative and positive powers of χ according to the elementary identities (157).

The elliptic solution (433) can also be written

$$u = s_1/3 + \partial_x \text{Log}(\text{ns}(\psi) - \text{cs}(\psi)), \quad \psi = \Psi_3, \quad g_3 = 0, \quad g_2 \text{ arb.}, \quad a_1 = 0, \quad (453)$$

a relation in which the argument of the logarithm is the ratio of two entire functions. Therefore it could be possible to find it by a suitable extension of a two-family truncation with a third order assumption.

This solution was first found by the following two-step procedure [10], which, unfortunately for this nice method, only works for a restricted class of PDEs (those with $p = -1$, $u_0 = c_0$, $u_1 = c_1C + c_2$, $c_j = \text{constant}$, see (80)). The first step is to define the truncation

$$\begin{aligned} u = d\partial_x \text{Log}(\varphi - \varphi_0) + U, \quad U = \text{constant}, \\ E(u) = \sum_{j=0}^3 E_j(\varphi_{xx}/\varphi_x, \varphi_t/\varphi_x) \left(\frac{\varphi - \varphi_0}{\varphi_x} \right)^{j-3}, \quad \forall j : E_j = 0, \end{aligned} \quad (454)$$

whose general solution is $U = s_1/3$, $\varphi - \varphi_0 = \Psi_3$ (indeed, comparing with (80), this assumption *a priori* implies $b\varphi_t - 3\varphi_{xx} = 0$). The second step is

not a truncation, but the change of function $u \mapsto f$

$$u = s_1/3 + (d\partial_x \text{Log } \Psi_3) f(\Psi_3), \quad (455)$$

which transforms (77) into

$$U'' - 2U^3 + 2a_1\Psi_{3,x}^{-3} = 0, \quad U(\psi) = f(\psi)/\psi. \quad (456)$$

This is an ODE iff $a_1 = 0$, in which case its solution is (433). Therefore, the assumption (455) has defined a reduction of the PDE to an ODE. This subject will be further examined in Section 10.

The cubic complex Ginzburg-Landau equation

The cubic complex Ginzburg-Landau equation (CGL3)

$$E(u) \equiv iu_t + pu_{xx} + q|u|^2u - i\gamma u = 0, \quad pq \neq 0, \quad (u, p, q) \in \mathcal{C}, \quad \gamma \in \mathcal{R}, \quad (457)$$

with p, q, γ constant, is a generic PDE describing the propagation of the signal in an optical fiber as well as superfluidity, spatiotemporal intermittency, pattern formation, etc.

One easily checks that $|u|$ generically behaves like a simple pole. The dominant behaviour

$$u \sim a_0\chi^{-1+i\alpha}, \quad \bar{u} \sim \bar{a}_0\chi^{-1-i\alpha}, \quad (458)$$

in which a_0 is a complex constant, α a real constant, is solution of the non-linear algebraic system

$$p(-1+i\alpha)(-2+i\alpha) + qa_2 = 0, \quad (459)$$

$$\bar{p}(-1-i\alpha)(-2-i\alpha) + \bar{q}a_2 = 0, \quad (460)$$

with $a_2 = |a_0|^2$. This defines two families for $|u|^2$ (four for $|u|$) [9]

$$a_2 = \frac{9|p|^2}{2|q|^2d_i^2}[d_r + \Delta], \quad \alpha = \frac{3}{2d_i}(d_r + \Delta), \quad (461)$$

$$\frac{p}{q} = d_r - id_i, \quad \Delta^2 = d_r^2 + (8/9)d_i^2. \quad (462)$$

To prevent these irrational expressions to mess up all subsequent computations (Fuchs indices, no-log conditions, truncations), the system (459)–(460) can equivalently be solved as a *linear* system on \mathcal{C} [31,35]

$$a_2 = -\frac{p}{q}(1-i\alpha)(2-i\alpha), \quad (463)$$

$$\bar{p} = Kp(1-i\alpha)(2-i\alpha), \quad \bar{q} = Kq(1+i\alpha)(2+i\alpha), \quad (464)$$

in which K is an irrelevant arbitrary nonzero complex constant.

The indicial equation is the determinant [26] of the second order matrix

$$\mathbf{P}(j) = \begin{pmatrix} (2a_0\overline{a_0})q & a_0^2q \\ \overline{q}a_0^2 & (2a_0\overline{a_0})\overline{q} \end{pmatrix} + \text{diag}(p(j-1+i\alpha)(j-2+i\alpha), \overline{p}(j-1-i\alpha)(j-2-i\alpha)), \quad (465)$$

and with the resolution (463)–(464) it evaluates to

$$\det \mathbf{P}(j) = (j+1)j(j^2 - 7j + 6\alpha^2 + 12) = 0. \quad (466)$$

For generic values of (p, q) , two of the four indices are irrational.

Let us consider, for simplification, the solitary wave reduction

$$u(x, t) = U(\xi)e^{i(\omega t + \varphi(\xi))}, \quad \xi = x - ct, \quad (467)$$

in which (U, φ) are functions of the reduced independent variable ξ , and let us restrict to the pure CGL3 case $\text{Im}(p/q) \neq 0$. The general solution of the fourth order system of ODEs for (U, φ) *a priori* depends on six arbitrary constants, the four constants of integration plus the two reduction parameters (c, ω) . From these six constants, one must subtract

1. the irrelevant origin ξ_0 of ξ (Fuchs index -1), which represents the invariance under a space translation,
2. the irrelevant origin φ_0 of the phase (Fuchs index 0), which represents the invariance under a phase shift,
3. and the number of irrational Fuchs indices, generically two. Indeed, these irrational indices represent the chaotic nature of CGL3 (see the expansion (48) in [31]) and they cannot contribute to any analytic solution.

Therefore only two relevant arbitrary constants are present in what can be called the *general analytic solution* of the reduction (467).

Presently, one only knows four particular solutions of the reduction $\xi = x - ct$ with a zero codimension (no constraint on (p, q, γ)). These are

1. a pulse or solitary wave [99]

$$u = -ia_0k \text{sech } kxe^{i[\alpha \text{Log} \cosh kx + K_1t]}, \quad K_2k^2 - \gamma = 0, \quad (468)$$

2. a front or shock [95]

$$u = a_0\frac{k}{2} \left[\tanh \frac{k}{2}\xi \pm 1 \right] e^{i[\alpha \text{Log} \cosh \frac{k}{2}\xi + K_3c\xi - K_4c^2t]}, \quad (469)$$

3. a source or propagating hole [5]

$$u = a_0 \left[\frac{k}{2} \tanh \frac{k}{2}\xi + (K_1 + iK_2)c \right] \times e^{i[\alpha \text{Log} \cosh \frac{k}{2}\xi + K_3c\xi - (K_4k^2 + K_5c^2)t]}, \quad K_6k^2 + K_7c^2 = \gamma, \quad (470)$$

4. an unbounded solution [31]

$$|u|^2 = a_2(\tan^2 \frac{k}{2}\xi + K^2), \quad c = 0. \quad (471)$$

In the above expressions, all parameters (a_2, α, k, c, K_l) are real and only depend on (p, q, γ) , except in (470) where the velocity c is arbitrary.

These four particular solutions are four different degeneracies [31] of the yet unknown general analytic solution.

In experiments or computer simulations, one has observed [5,73,12,56] other regular patterns which should correspond to other degeneracies of the general analytic solution. One of them [56] is a homoclinic hole solution, complementing the heteroclinic hole (470). Another one, of the highest interest in fiber optics, is a propagating pulse, extrapolating (468) to $c \neq 0$ and reducing in the NLS limit (p, q real, $\gamma = 0$) to a “bright soliton” of arbitrary velocity.

Let us now address the question of retrieving these four solutions (and ideally of finding the unknown one ar at least other degeneracies) by some truncation. For a truncation to be successful, the truncated variables should be free of any multivaluedness in their dominant behaviour. This is not the case of the natural physical variables (u, \bar{u}) or $(\operatorname{Re} u, \operatorname{Im} u)$, which are *always* locally multivalued as seen from (458). A more detailed study [31] uncovers the best representation for this purpose, namely a *complex modulus* Z and a real argument Θ uniquely defined by

$$u = Ze^{i\Theta}, \quad \bar{u} = \bar{Z}e^{-i\Theta}, \quad (472)$$

and the above four exact solutions are written in this notation. For each family, if one excludes the contribution of the irrational Fuchs indices, the three fields $(Z, \bar{Z}, \operatorname{grad} \Theta)$ are locally singlevalued and they behave like simple poles. The physical variables $(|u|^2, \operatorname{grad} \arg u)$ also have this nice property of being locally singlevalued (they respectively behave like a double pole and a simple pole), but they are not as elementary as $(Z, \bar{Z}, \operatorname{grad} \Theta)$.

The one-family truncation of the third order ODE satisfied by $|u|^2$ (after elimination of φ), with a constant coefficient second order assumption, evidently captures all four solutions, since $|u|^2$ is a degree-two polynomial in $\tanh \kappa \xi$. Such a truncation generates cumbersome computations and provides no additional solution.

The one-family truncation of $(Z, \bar{Z}, \operatorname{grad} \Theta)$ with the same constant coefficient second order assumption is defined as [31,33]

$$\begin{cases} Z = a_0(\chi^{-1} + X + iY), \\ \bar{Z} = \bar{a}_0(\chi^{-1} + X - iY), \\ \Theta = \omega t + \alpha \operatorname{Log} \psi + K\xi, \\ (\operatorname{Log} \psi)' = \chi^{-1}, \quad \chi' = 1 - (k^2/4)\chi^2, \\ Ee^{-i\Theta} = \sum_{j=0}^3 E_j \chi^{j-3}, \end{cases} \quad (473)$$

in which χ and ψ are functions of $\xi = x - ct$, (ω, X, Y, K, k^2) are real constants. One has to solve the four complex (eight real) equations $E_j = 0$ in the eight

real unknowns $(a_2, \alpha, \omega, X, Y, K, c, k^2)$, the two complex parameters (p, q) , and the real parameter γ . If there exists a solution, the elementary building block functions evaluate to

$$\chi = \frac{k}{2} \tanh \frac{k\xi}{2}, \quad \psi = \cosh \frac{k\xi}{2}. \quad (474)$$

The good methodology [31,35] is again to select, among the eleven complex variables considered as equivalent, four variables which make the system a *linear* one of Cramer type. The system (E_0, E_1, E_2) is of Cramer type in (a_2, K, ω) , and after its resolution the last equation E_3 is independent of (p, q, γ, c) and factorizes into a product of *linear* factors

$$E_3 \equiv [k^2 - 4(X + iY)^2](\alpha Y - 2X) = 0. \quad (475)$$

Finally, this one-family truncation recovers all four solutions except the pulse (468).

The two-family truncation of $(Z, \bar{Z}, \text{grad } \Theta)$ with the same constant coefficient second order assumption retrieves the pulse solution (468), but finds nothing new.

Similar truncations for two coupled CGL3 equations can be found in [35,36].

The nonintegrable Kundu-Eckhaus equation

The PDE for the complex field $U(x, t)$ [68,8]

$$iU_t + \alpha U_{xx} + \left(\frac{\beta^2}{\alpha} |U|^4 + 2be^{i\gamma}(|U|^2)_x\right)U = 0, \quad (\alpha, \beta, b, \gamma) \in \mathcal{R}, \quad (476)$$

with $\alpha\beta \cos \gamma \neq 0$, is linearizable when $b^2 = \beta^2$ into the Schrödinger equation

$$iV_t + \alpha V_{xx} = 0, \quad U = \sqrt{\frac{\alpha}{2\beta \cos \gamma}} \frac{V}{\sqrt{\int |V|^2 dx}}. \quad (477)$$

This suggests considering the PDE for $u = \int |V|^2 dx$

$$\begin{aligned} & \frac{\alpha}{2}(u_{xxxx}u_x^2 + u_{xx}^3 - 2u_x u_{xx} u_{xxx}) + 2\frac{\beta^2 - (b \sin \gamma)^2}{\alpha} u_x^4 u_{xx} \\ & + 2(b \cos \gamma) u_x^3 u_{xxx} + \frac{1}{2\alpha}(u_{tt}u_x^2 + u_{xx}u_t^2 - 2u_t u_x u_{xt}) = 0. \end{aligned} \quad (478)$$

When $b^2 \neq \beta^2$, this PDE fails the test [15] because, for each of the two families for u , one index is generically irrational. However, its one-family truncation with a second order assumption (i.e. the usual WTC truncation) is a very rich exercise [32] which yields quite unusual solutions, among them an elliptic one involving the ODE of class III of Chazy [13].

10 Singular manifold method *versus* reduction methods

In order to find exact solutions of PDEs, there exist two main classes of methods. The first class, which has been detailed in these lectures, is based on the structure of the movable singularities and it can be called, in short, the singular manifold method.

The second class, presented in another course of this school [121], basically relies on group theory and consists in finding the *reductions* to a PDE in a lesser number of independent variables, and at the end to an ODE. These reductions are obtained either by looking for the infinitesimal symmetries of the PDE (space translation, etc) and by integrating them, or by a direct search not involving any group theory. The main methods in this second class are known as (see references in [121]) the *classical method* (point symmetries), the *nonclassical method* (conditional point symmetries), the *direct method* (direct search).

The question of the comparison of these four methods by their results is an active research subject [44,45,52], and its current state is given in [17,121].

Let us take as an example a second order nonintegrable PDE, this is enough to give an idea of the comparison. The KPP equation (77) has been studied in detail with the singular manifold method, Section 9.3. It has also been investigated with the three other methods, and the results are the following.

In the classical and nonclassical methods, let us denote

$$\tau(x, t, u)u_t + \xi(x, t, u)u_x - \eta(x, t, u) = 0 \quad (479)$$

the PDE for $u(x, t)$ which, after computation of the symmetries (τ, ξ, η) , defines the constraint on u susceptible to yield a reduction if the constraint can be integrated.

Classical method. It yields only two reductions $u(x, t) \mapsto U(z)$ [9,10], one noncharacteristic (i.e. conserving the differential order two)

$$z = x - ct, \quad u = U, \quad -U'' - bcU' + 2d^{-2}(U - e_1)(U - e_2)(U - e_3) = 0, \quad (480)$$

one characteristic (i.e. lowering the differential order two)

$$z = t, \quad u = U, \quad bU' + 2d^{-2}(U - e_1)(U - e_2)(U - e_3) = 0. \quad (481)$$

This is in fact a unique reduction $z = \lambda x + \mu t$, but the splitting according to the characteristic nature is relevant for the Painlevé property. None of these two ODEs has the Painlevé property, unless $c = (3e_j - s_1)/(bd)$ in (480).

Nonclassical method. It yields three sets of values for (τ, ξ, η) [96], two with $\tau \neq 0$ and one with $\tau = 0, \xi \neq 0$.

The first one [10] has codimension one

$$a_1 = 0, \quad e_1 = (e_2 + e_3)/2, \quad bu_t - 3d^{-2}((\text{Log } \psi)_x(u - e_1))_x = 0, \quad \psi = \Psi_3, \quad (482)$$

its integration defines the noncharacteristic reduction to an elliptic equation

$$a_1 = 0, \quad z = \Psi_3, \quad u = e_1 + dz_x U(z), \quad U'' - 2U^3 = 0, \quad (483)$$

and one finds the solution (433).

The second one [96] has codimension zero

$$bu_t + d^{-1}(u - s_1/3)u_x + 3d^{-2}(u - e_1)(u - e_2)(u - e_3) = 0, \quad (484)$$

and, remarkably, this first order PDE, which fails the test, identifies to the no-log condition (82) with $3u - s_1 = bdC$. Its integration (436) cannot define a reduction unless some choice of the arbitrary function is made. Nevertheless, the common solution to the PDEs (77) and (484) is (431).

The third one [96] is

$$u_x - \eta = 0, \quad (485)$$

in which η satisfies the second order PDE

$$\begin{aligned} \eta_{xx} + 2\eta\eta_{xu} + \eta^2\eta_{uu} + 2d^{-2}(u - e_1)(u - e_2)(u - e_3)\eta_u \\ - 2d^{-2}(3u^2 - 2s_1u + s_1^2/3 - 3d^2a_2)\eta + b\eta_t = 0. \end{aligned} \quad (486)$$

Integrating (486) is equivalent to integrating the original PDE (77), since the transformation (485) simply exchanges them, so one is stranded. The only way out is to put some additional constraints on η . The consistent way to do that ([17] page 634) is to eliminate u_x and its derivatives (in this case u_{xx} only) between (485) and (77), which results in a nonlinear first order ODE for the function $t \mapsto u(x, t)$ (i.e. with x as a parameter)

$$bu_t - (\eta_x + \eta\eta_u) + 2d^{-2}(u - e_1)(u - e_2)(u - e_3) = 0. \quad (487)$$

Requiring the invariance of this ODE under the infinitesimal transformation ($\tau = 0, \xi = 1, \eta$) of the classical method creates constraints on η , an exercise which is left to the reader.

Direct method. The search for a reduction $u(x, t) \mapsto U(z)$ in the class

$$u(x, t) = \lambda(x, t)U(z(x, t)) + \mu(x, t), \quad (488)$$

apart from the characteristic reduction $z = t, u = U$, yields the noncharacteristic reduction [96]

$$U'' - 2U^3 + g_1(z)U' + g_2(z)U + g_3(z) = 0, \quad (489)$$

$$u = dz_x U + s_1/3, \quad z_x \neq 0, \quad (490)$$

provided z, g_1, g_2, g_3 satisfy the system

$$\begin{cases} bz_{xt} - z_{xxx} - 6a_2z_x + z_x^3g_2 = 0, \\ bz_t - 3z_{xx} + z_x^2g_1 = 0, \\ 2a_1 - z_x^3g_3 = 0. \end{cases} \quad (491)$$

The constraint $z_x \neq 0$ splits the discussion into $a_1 = 0$ and $a_1 \neq 0$. The case a_1 arbitrary defines the reduction [96]

$$z = x - ct, \quad u = dU + s_1/3, \quad U'' + cU' - 2(U^3 - 3a_2U - a_1) = 0, \quad (492)$$

identical to (480). In the case $a_1 = 0$, hence $g_3 = 0$, the system is solved for (z_{xxx}, z_t) , and the condition $(z_{xxx})_t = (z_t)_{xxx}$ reads

$$\left(18 \left(\frac{z_{xx}}{z_x^2} \right)^2 + 18 \frac{z_{xx}}{z_x^2} + 2g_1'(z) - g_1(z)\partial_z + 3\partial_z^2 \right) Q(z) = 0, \quad (493)$$

$$Q(z) = 9g_2 - 2g_1^2 - 3g_1'. \quad (494)$$

For $Q(z) \neq 0$, the condition integrates as

$$z = G(xf_1(t) + f_2(t)), \quad (495)$$

in which G is an arbitrary function, and (f_1, f_2) are further constrained by $f_1' = 0, f_2' = 0$. The result is $a_1 = 0, z = G(x)$, and the ODE (489) transforms to an elliptic equation under

$$U(z) = \frac{f(G^{-1}(z))}{G'(z)}, \quad f'' - 2f^3 + 6a_2f = 0, \quad (496)$$

in which G^{-1} denotes the inverse function of G . This solution is not distinct from the stationary elliptic reduction (434).

For $Q(z) = 0$, if one defines the function G by

$$g_1(z) = -3(\text{Log}(G'(z)))', \quad (497)$$

the system (491) is equivalent to

$$\begin{cases} g_2 = (2/9)g_1^2 + (1/3)g_1', \\ (\partial_x^3 - 3a_2\partial_x)G(z(x, t)) = 0, \\ (b\partial_t - 3\partial_x^2)G(z(x, t)) = 0, \\ f''(Z) - 2f(Z)^3 = 0, \quad U(z) = G'(z)f(Z), \quad Z = G(z), \end{cases} \quad (498)$$

and this proves that the particular solution $g_1 = g_2 = 0$ considered in [96] is the general solution, equivalent to the reduction $z = \Psi_3$ in (433).

11 Truncation of the unknown, not of the equation

When applied for instance to the second Painlevé equation (P2)

$$(P2) \quad E(u) \equiv u'' - 2u^3 - xu - \alpha = 0, \quad (499)$$

the one-family singular manifold method in the case of a second order scattering problem, i.e. the one originally performed by WTC, see Sections 7.1

and 7.3, presents the following drawback. The (P2) ODE has two families $u \sim \varepsilon\chi^{-1}$, $\varepsilon^2 = 1$. With the definition of $\text{grad } \chi$

$$\chi' = 1 + (S/2)\chi^2, \quad (500)$$

the one-family truncated expansion for u is found to be

$$u = \varepsilon\chi^{-1}, \quad E(u) = \varepsilon(S - x)\chi^{-1} + (-\alpha - \varepsilon S'/2)\chi^0, \quad (501)$$

$$E_2 \equiv \varepsilon(S - x) = 0, \quad (502)$$

$$E_3 \equiv -\alpha - \varepsilon S'/2 = 0, \quad (503)$$

and its general solution is

$$S = x, \quad 2\alpha + \varepsilon = 0, \quad u = \varepsilon(\text{Log Ai}(x))', \quad \text{Ai}'' + (x/2)\text{Ai} = 0. \quad (504)$$

One therefore finds only a one-parameter particular solution in terms of the Airy function, at the price of one constraint on the parameter α . This is unsatisfactory because the method fails to find the highest information on (P2) (highest in the context of these lectures), namely its *Schlesinger transformation*. Such a transformation is by definition a *birational transformation* between two different copies of (P2), denoted $u(x, \alpha)$ and $U(X, A)$, and it reads [75]

$$x = X, \quad u + U = \frac{-2A - 1}{2(U' + U^2) + X} = \frac{2\alpha - 1}{2(u' - u^2) - x}, \quad \alpha = A + 1. \quad (505)$$

A method to remedy this drawback is the following [16]. We rephrase it in the homographically invariant formalism, which simplifies the exposition. Firstly, rather than splitting $E(u)$, defined in (501), into one equation per power of χ , one retains the single information $E(u) = 0$, and one eliminates u and χ between the three equations (500) and (501) to obtain the second order ODE for $S(x)$

$$2(S - x)S'' - S'^2 + 2S' + 2S^3 - 4xS^2 + 2x^2S + 4\alpha(\alpha + \varepsilon) = 0. \quad (506)$$

This ODE for $S(x)$, which is birationally equivalent to (P2) under the transformation

$$u = \varepsilon\chi^{-1}, \quad S = -2(\chi^{-1})' - 2\chi^{-2}, \quad \chi^{-1} = \frac{S' + 2\varepsilon\alpha}{2(S - x)}, \quad (507)$$

bears the number 34 in the classification of Painlevé and Gambier [51].

Secondly, despite the fact that one already knows the general solution $S(x)$ in terms of the (P2) function $u(x, \alpha)$, one takes advantage of the two-family structure of (P2) (the sign ε is ± 1) to perform an involution by representing $S(x)$ with another (P2) function $U(X, A)$ as

$$S = -2V' - 2V^2, \quad U = \varepsilon_2 V, \quad \varepsilon_2^2 = 1, \quad U'' - 2U^3 - XU - A = 0, \quad X = x. \quad (508)$$

The elimination of S between (507) and (508) provides a relation between $(\varepsilon\alpha, \varepsilon_2A)$ only

$$(A + \varepsilon_2/2)^2 = (\alpha + \varepsilon/2)^2. \quad (509)$$

The solution $A = -\varepsilon_2(\varepsilon\alpha + 1)$ is the Schlesinger transformation.

An equivalent presentation can be found in Ref. [55]. In the latter, one first computes the two coefficients u_0, u_1 of the Laurent expansion

$$u = u_0\chi^{-1} + u_1, \quad (510)$$

then the Schlesinger transformation is readily obtained by (more precisely, the computation of [55] reduces to) the elimination of the three variables u, Z, U'' between the four equations (u_0, u_1, u, U, Z, s are functions of $X = x$)

$$u = u_0Z^{-1} + U, \quad (511)$$

$$Z' = 1 + 2\frac{U - u_1}{u_0}Z + \frac{s}{2}Z^2, \quad (512)$$

$$(\text{Pn})(u, x, \alpha, \beta, \gamma, \delta) = 0, \quad (513)$$

$$(\text{Pn})(U, X, A, B, C, D) = 0. \quad (514)$$

Equation (511) is an assumption for a Darboux transformation, and (512) defines a Riccati equation for the expansion variable Z which depends on a free function s . The elimination is differential for u and Z , algebraic for U'' , and it results in

$$F(U', U; s, s', s'', \alpha, \beta, \gamma, \delta, A, B, C, D) = 0. \quad (515)$$

The algebraic independence of (U', U) , consequence of the irreducibility of (Pn) , requires the identical vanishing of F as a polynomial of the two variables (U', U) , and this provides two solutions: the identity ($u = U, Z^{-1} = 0$) and, at least for (P2) and (P4), the Schlesinger transformation. The result for (P2) is

$$(\text{P2}) \quad \varepsilon Z^{-1} = u - U = \frac{\varepsilon(A - \alpha)}{2U' + \varepsilon(2U^2 + x)}, \quad \alpha + A + \varepsilon = 0, \quad s = 0, \quad (516)$$

and the inverse transformation

$$(\text{P2}) \quad u - U = \frac{\varepsilon(A - \alpha)}{2u' + \varepsilon(2u^2 + x)} \quad (517)$$

follows from the elimination of U' between (516) and

$$(U - u)' + \varepsilon(U^2 - u^2) = 0, \quad (518)$$

itself obtained by the elimination of Z between (511) and (512). This Schlesinger transformation is identical, thanks to the parity invariance of (P2), to (505).

The result for (P4) is

$$(P4) \quad u'' - u'^2/(2u) - (3/2)u^3 - 4xu^2 - 2x^2u + 2\alpha u - \beta/u = 0, \quad (519)$$

$$\begin{aligned} \varepsilon Z^{-1} = u - U &= \frac{4\varepsilon(\alpha - A)U}{3U' + \varepsilon(3U^2 + 6xU - 2A - 4\alpha) + 6}, \\ &= \frac{4\varepsilon(\alpha - A)u}{3u' + \varepsilon(3u^2 + 6xu - 2\alpha - 4A) + 6}, \end{aligned} \quad (520)$$

$$(U - u)' + \varepsilon(U^2 - u^2 + 2x(U - u) + 2(\alpha - A)/3) = 0, \quad (521)$$

$$9\beta + 2(\alpha + 2A - 3\varepsilon)^2 = 0, \quad 9B + 2(A + 2\alpha - 3\varepsilon)^2 = 0, \quad (522)$$

$$s = 4(A - \alpha)/3. \quad (523)$$

12 Conclusion, open problems

The singular manifold method, which is based on the singularity structure, is quite powerful to provide exact solutions or other analytic results. There still exist many challenging problems, in particular in nonlinear optics and spatiotemporal intermittency [5,56], in which the equations, although nonintegrable, possess some regular “patterns” which could well be described by exact particular solutions. The difficulty to find them [31] comes from the good guess which must be made for the functions ψ , which do not necessarily satisfy a linear system any more. Methods from group theory usually provide complementary results, although they also fail in the two just quoted examples.

Acknowledgments

The author thanks the CIME Fondazione and l’Università di Palermo for invitation.

Index

- N -soliton solution, 4
- ψ -series, 14, 49

- AKNS equation, 32, 60
- AKNS system, 58, 60

- Bäcklund parameter, 4, 5, 26
- Bäcklund transformation, 3, 19, 48
 - auto-, 4, 58
 - bilinear, 48, 59
 - hetero-, 4, 5, 11, 26, 31, 34, 36, 44, 45, 58
- birational transformation, 71
- Boussinesq equation, 20, 35, 53
 - modified, 59
- branch, 15
- Broer-Kaup equation, 20, 59, 60

- characteristic, 69
- classical
 - method, 68
- closed form, 3
- complex Ginzburg-Landau equation, 64
- Robert Conte, 1

- d'Alembert equation, 5, 11, 21, 30, 31
- Darboux transformation, 8, 19, 20
 - binary, 38
- differential Groebner basis, 50
- differential part, 7
- diophantine conditions, 17
- direct
 - method, 68

- elliptic function, 18
- expansion variable, 12

- family of movable singularities, 15
- Fordey-Gibbons equation, 44
- Fuchs indices, 15

- general analytic solution, 52, 65

- hierarchy, 20
- Hirota-Satsuma equation, 20, 37, 41, 53

- homographic
 - group, 13
 - invariants, 13

- indicial equation, 15
- integrability, 5
- isospectral, 6
- Ito equation, 8

- Kaup-Kupershmidt equation, 20, 42–45, 53, 60
- Korteweg-de Vries
 - hierarchy, 8
- Korteweg-de Vries equation, 12, 18, 33, 58, 60
 - modified, 57, 60
- KPP equation, 16, 60, 68
- Kundu-Eckhaus equation, 67
- Kuramoto-Sivashinsky equation, 3, 5, 9, 17, 18, 49

- Lax pair, 6, 19
 - Lax representation, 6
 - order, 6
 - Riccati representation, 6
 - scalar representation, 6
 - string representation, 6
 - zero-curvature representation, 6
- Liouville equation, 5, 11, 30, 55, 57, 60
- Lorenz model, 18

- microdifferential operator, 7
- Miura transformation, 35

- no-logarithm condition, 15
- nonclassical
 - method, 68
- nonlinear Schrödinger equation, 7, 58, 60, 66

- one-soliton solution, 4

- Painlevé
 - analysis, 11
 - equation (P1), 8

- equation (P2), 71, 72
- equation (P4), 72
- equations, 19
- property, 10
- test, 11

Rössler system, 18

reduction, 61, 64, 65, 68, 69

resonances, 15

resummation, 20, 21

Sawada-Kotera equation, 20, 42, 43, 45,
53, 60

Schlesinger transformation, 19

Schwarzian, 13

sine-Gordon equation, 4, 5, 7, 9, 11, 18,
20, 55, 60

singular manifold

- equation, 29, 33, 35, 36, 51, 53, 56,
58

- method, 19, 20, 27, 68

- variable, 11

singular part operator, 16

solution

- exact, 3, 18

- global, 3

- local, 3

spectral parameter, 6, 26

tau-function, 8, 19

truncation

- equations, 27

- method, 19, 20, 27

- variable, 27

Tzitzéica equation, 20, 38, 60

vacuum, 4

References

1. M. J. Ablowitz and P. A. Clarkson, *Solitons, nonlinear evolution equations and inverse scattering* (Cambridge University Press, Cambridge, 1991).
2. M. J. Ablowitz, D. J. Kaup, A. C. Newell and H. Segur, The inverse scattering transform-Fourier analysis for nonlinear problems, *Stud. Appl. Math.* **53** (1974) 249–315.
3. I. Ayari and A. M. Grundland, Conditional symmetries for k th order partial differential equations, 24 pages, preprint CRM–2488 (1997).
4. A. V. Bäcklund, Om ytor med konstant negativ krökning, *Lunds Universitets Arsskrift Avd. 2* **19** (1883).
5. N. Bekki and K. Nozaki, Formations of spatial patterns and holes in the generalized Ginzburg-Landau equation, *Phys. Lett. A* **110** (1985) 133–135.
6. F. Boulier, *Étude et implantation de quelques algorithmes en algèbre différentielle* (Thèse, Université des sciences et technologies de Lille, 1994).
7. S. Bouquet, Hamiltonian structure and integrability of the stationary Kuramoto-Sivashinsky equation, *J. Math. Phys.* **36** (1995) 1242–1258.
8. F. Calogero and W. Eckhaus, Nonlinear evolution equations, rescalings, model PDEs and their integrability, I, *Inverse problems* **3** (1987) 229–262.
9. F. Cariello and M. Tabor, Painlevé expansions for nonintegrable evolution equations, *Physica D* **39** (1989) 77–94.
10. F. Cariello and M. Tabor, Similarity reductions from extended Painlevé expansions for nonintegrable evolution equations, *Physica D* **53** (1991) 59–70.
11. P. J. Caudrey, The inverse problem for the third order equation $u_{xxx} + q(x)u_x + r(x)u = -i\zeta^3 u$, *Phys. Lett. A* **79** (1980) 264–268.
12. H. Chaté, Spatiotemporal intermittency régimes of the one-dimensional complex Ginzburg-Landau equation, *Nonlinearity* **7** (1994) 185–204.
13. J. Chazy, Sur les équations différentielles du troisième ordre et d'ordre supérieur dont l'intégrale générale a ses points critiques fixes, Thèse, Paris (1910); *Acta Math.* **34** (1911) 317–385.
14. H. H. Chen, General derivation of Bäcklund transformations from inverse scattering problems, *Phys. Rev. Lett.* **33** (1974) 925–928.
15. P. A. Clarkson and C. M. Cosgrove, The Painlevé property and a generalised derivative nonlinear Schrödinger equation, *J. Phys. A* **20** (1987) 2003–2024.
16. P. A. Clarkson, N. Joshi, and A. Pickering, Bäcklund transformations for the second Painlevé hierarchy: a modified truncation approach, *Inverse Problems* **15** (1999) 175–187.
17. P. A. Clarkson and P. Winternitz, Symmetry reduction and exact solutions of nonlinear partial differential equations, *The Painlevé property, one century later*, 591–660, ed. R. Conte, CRM series in mathematical physics (Springer, New York, 1999).
18. R. Conte, Universal invariance properties of Painlevé analysis and Bäcklund transformation in nonlinear partial differential equations, *Phys. Lett. A* **134** (1988) 100–104.
19. R. Conte, Invariant Painlevé analysis of partial differential equations, *Phys. Lett. A* **140** (1989) 383–390.
20. R. Conte, Painlevé singular manifold equation and integrability, *Inverse methods in action*, pp. 497–504, ed. P. C. Sabatier, Springer-Verlag series “Inverse problems and theoretical imaging” (Springer-Verlag, Berlin, 1990).

21. R. Conte, Painlevé analysis and Bäcklund transformation of the nonintegrable KPP equation, *Nonlinear evolution equations: integrability and spectral methods*, 187–192, eds. A. Degasperis, A. P. Fordy, and M. Lakshmanan (Manchester University Press, Manchester, 1990).
22. R. Conte, Towards the equivalence between integrability and Painlevé test for partial differential equations, *Nonlinear and chaotic phenomena in plasmas, solids and fluids*, 94–101, eds. W. Rozmus and J. A. Tuszynski (World Scientific, Singapore, 1991).
23. R. Conte, Unification of PDE and ODE versions of Painlevé analysis into a single invariant version, *Painlevé transcendents, their asymptotics and physical applications*, 125–144, eds. D. Levi and P. Winternitz (Plenum, New York, 1992).
24. R. Conte, Exact solutions of nonlinear wave equations by singularity methods, *Rendiconti del Circolo Matematico di Palermo*, Suppl. **57** (1998) 165–175.
25. R. Conte (ed.), *The Painlevé property, one century later*, 810 pages, CRM series in mathematical physics (Springer, New York, 1999).
26. R. Conte, The Painlevé approach to nonlinear ordinary differential equations, *The Painlevé property, one century later*, 77–180, ed. R. Conte, CRM series in mathematical physics (Springer, New York, 1999). solv-int/9710020.
27. R. Conte, Various truncations in Painlevé analysis of partial differential equations, 103–116, *Nonlinear dynamics : integrability and chaos*, eds. M. Daniel, K. M. Tamizhmani, and R. Sahadevan (Narosa publishing house, New Delhi, 2000). solv-int/9812008.
28. R. Conte, A. P. Fordy and A. Pickering, A perturbative Painlevé approach to nonlinear differential equations, *Physica D* **69** (1993) 33–58.
29. R. Conte and M. Musette, Painlevé analysis and Bäcklund transformation in the Kuramoto-Sivashinsky equation, *J. Phys. A* **22** (1989) 169–177.
30. R. Conte and M. Musette, Link between solitary waves and projective Riccati equations, *J. Phys. A* **25** (1992) 5609–5623.
31. R. Conte and M. Musette, Linearity inside nonlinearity: exact solutions to the complex Ginzburg-Landau equation, *Physica D* **69** (1993) 1–17.
32. R. Conte and M. Musette, Exact solutions to the partially integrable Eckhaus equation, *Nonlinear evolution equations and dynamical systems*, eds. L. Martina and A. K. Pogrebkov, *Teoreticheskaya i Matematicheskaya Fizika* **99** (1994) 226–233 [English : *Theor. and Math. Phys.* **99** (1994) 543–548].
33. R. Conte and M. Musette, Exact solutions to the complex Ginzburg-Landau equation of nonlinear optics, *Pure Appl. Opt.* **4** (1995) 315–320.
34. R. Conte and M. Musette, Beyond the two-singular manifold method, *Nonlinear physics : theory and experiment*, 67–74, eds. E. Alfinito, M. Boiti, L. Martina and F. Pempinelli (World Scientific, Singapore, 1996).
35. R. Conte and M. Musette, Analytic expressions of hydrothermal waves, 12 pages, Reports on mathematical physics, to appear (2000). Preprint S99/054. nlin-si/0009022.
36. R. Conte and M. Musette, On the solitary wave of two coupled nonintegrable Ginzburg-Landau equations, *Nonlinear integrability and all that : twenty years after NEEDS '99*, 382–388, eds. M. Boiti, L. Martina, F. Pempinelli, B. Prinari, and G. Soliani (World Scientific, Singapore, 2000). Preprint S99/058.
37. R. Conte, M. Musette and A. M. Grundland, Bäcklund transformation of partial differential equations obtained from the Painlevé-Gambier classification, II. Tzitzéica equation, *J. Math. Phys.* **40** (1999) 2092–2106.

38. R. Conte, M. Musette and A. Pickering, The two-singular manifold method, II. Classical Boussinesq system, *J. Phys. A* **28** (1995) 179–185.
39. G. Contopoulos, B. Grammaticos and A. Ramani, Painlevé analysis for the mixmaster universe model, *J. Phys. A* **25** (1993) 5795–5799.
40. G. Darboux, Sur une proposition relative aux équations linéaires, *C. R. Acad. Sc. Paris* **94** (1882) 1456–1459.
41. G. Darboux, *Leçons sur la théorie générale des surfaces et les applications géométriques du calcul infinitésimal*, vol. III (Gauthier-Villars, Paris, 1894). Reprinted, *Théorie générale des surfaces* (Chelsea, New York, 1972). Reprinted (Gabay, Paris, 1993).
42. P. Di Francesco, 2-D quantum and topological gravities, matrix models, and integrable differential systems, *The Painlevé property, one century later*, 229–285, ed. R. Conte, CRM series in mathematical physics (Springer, New York, 1999).
43. P. G. Estévez, P. R. Gordoa, L. Martínez Alonso, and E. Medina Reus, Modified singular manifold expansion : application to the Boussinesq and Mikhailov-Shabat systems, *J. Phys. A* **26** (1993) 1915–1925.
44. P. G. Estévez and P. R. Gordoa, Nonclassical symmetries and the singular manifold method : theory and six examples, *Stud. Appl. Math.* **95** (1995) 73–113.
45. P. G. Estévez and P. R. Gordoa, Nonclassical symmetries and the singular manifold method : a further two examples, *J. Phys. A* **31** (1998) 7511–7519.
46. A. P. Fordy and J. Gibbons, Some remarkable nonlinear transformations, *Phys. Lett. A* **75** (1980) 325–325.
47. A. P. Fordy and A. Pickering, *Symmetries and Integrability of Difference equations* London Mathematical Society LNS **255** (1999) 287–298.
48. A. R. Forsyth, *Theory of differential equations*. Part IV (vol. VI) Partial differential equations (Cambridge University Press, Cambridge, 1906). Reprinted (Dover, New York, 1959).
49. J.-D. Fournier, E. A. Spiegel and O. Thual, Meromorphic integrals of two non-integrable systems, *Nonlinear dynamics*, 366–?, eds. G. Servizi and G. Turchetti (World Scientific, Singapore, 1989).
50. B. Gaffet, A class of 1–d gas flows soluble by the inverse scattering transform, *Physica D* **26** (1987) 123–139.
51. B. Gambier, Sur les équations différentielles du second ordre et du premier degré dont l’intégrale générale est à points critiques fixes, Thèse, Paris (1909); *Acta Math.* **33** (1910) 1–55.
52. M.L. Gandarias and M.S. Bruzón, Nonclassical symmetries for a family of Cahn-Hilliard equations, *Phys. Lett. A* **263** (2000) 331–337.
53. T. I. Garagash, On a modification of the Painlevé test for the BLP equation, *Nonlinear evolution equations and dynamical systems*, 130–133, eds. V. G. Makhankov, I. V. Puzynin, and O. K. Pashaev (World Scientific, Singapore, 1993).
54. J. D. Gibbon, P. Radmore, M. Tabor, and D. Wood, The Painlevé property and Hirota’s method, *Stud. Appl. Math.* **72** (1985) 39–63.
55. P. Gordoa, N. Joshi, and A. Pickering, Mappings preserving locations of movable poles: a new extension of the truncation method to ordinary differential equations, *Nonlinearity* **12** (1999) 955–968.
56. M. van Hecke, Building blocks of spatiotemporal intermittency, *Phys. Rev. Lett.* **80** (1998) 1896–1899.

57. E. Hille, *Ordinary differential equations in the complex domain* (J. Wiley and sons, New York, 1976).
58. R. Hirota, Fundamental properties of the binary operators in soliton theory and their generalization, *Dynamical problems in soliton systems*, 42–49, ed. S. Takeno (Springer, Berlin, 1985)
59. R. Hirota and A. Ramani, The Miura transformations of Kaup's equation and of Mikhailov's equation, *Phys. Lett. A* **76** (1980) 95–96.
60. R. Hirota and J. Satsuma, N-soliton solutions of model equations for shallow water waves, *J. Phys. Soc. Japan Letters* **40** (1976) 611–612.
61. R. Hirota and J. Satsuma, Nonlinear evolution equations generated from the Bäcklund transformation for the Boussinesq equation, *Prog. Theor. Phys.* **57** (1977) 797–807.
62. M. Ito, An extension of nonlinear evolution equations of the K-dV (mK-dV) type to higher orders, *J. Phys. Soc. Japan* **49** (1980) 771–778.
63. M. Jimbo, M. D. Kruskal and T. Miwa, Painlevé test for the self-dual Yang-Mills equation, *Phys. Lett. A* **92** (1982) 59–60.
64. M. Jimbo and T. Miwa, Solitons and infinite dimensional Lie algebras, *Publ. RIMS, Kyoto* **19** (1983) 943–1001.
65. D. J. Kaup, On the inverse scattering problem for cubic eigenvalue problems of the class $\psi_{xxx} + 6Q\psi_x + 6R\psi = \lambda\psi$, *Stud. Appl. Math.* **62** (1980) 189–216.
66. T. Kawahara and M. Tanaka, Interactions of traveling fronts : an exact solution of a nonlinear diffusion equation, *Phys. Lett. A* **97** (1983) 311–314.
67. A. N. Kolmogorov, I. G. Petrovskii and N. S. Piskunov, The study of a diffusion equation, related to the increase of the quantity of matter, and its application to one biological problem, *Bulletin de l'Université d'État de Moscou, série internationale, section A Math. Méc.* **1** (1937) 1–26.
68. A. Kundu, Landau-Lifshitz and higher order nonlinear systems gauge generated from nonlinear Schrödinger type equations, *J. Math. Phys.* **25** (1984) 3433–3438.
69. Y. Kuramoto and T. Tsuzuki, Persistent propagation of concentration waves in dissipative media far from thermal equilibrium, *Prog. Theor. Phys.* **55** (1976) 356–369.
70. G. L. Lamb Jr, Propagation of ultrashort optical pulses, *Phys. Lett. A* **25** (1967) 181–182.
71. G. L. Lamb Jr, Bäcklund transformations for certain nonlinear evolution equations, *J. Math. Phys.* **15** (1974) 2157–2165.
72. P. D. Lax, Integrals of nonlinear equations of evolution and solitary waves, *Comm. Pure Appl. Math.* **21** (1968) 467–490.
73. J. Lega, B. Jانياud, S. Jucquois and V. Croquette, Localized phase jumps in wave trains *Phys. Rev. A* **45** (1992) 5596–5604.
74. D. Levi and O. Ragnisco, Non-isospectral deformations and Darboux transformations for the third-order spectral problem, *Inverse Problems* **4** (1988) 815–828.
75. N. A. Lukashovich, The second Painlevé equation, *Differentsial'nye Uravneniya* **7** (1971) 1124–1125 [English : *Diff. equ.* **7** (1971) 853–854].
76. F. Magri, Chapter ?, this volume.
77. V. G. Makhankov and O. K. Pashaev, Nonlinear Schrödinger equation with noncompact isogroup, *Teoreticheskaya i Matematicheskaya Fizika* **53** (1982) 55–67 [English : *Theor. and Math. Phys.* **53** (1982) 979–987].

78. P. Manneville, The Kuramoto-Sivashinsky equation : a progress report, *Propagation in systems far from equilibrium*, 265–280, eds. J. Weisfreid, H. R. Brand, P. Manneville, G. Albinet, and N. Boccara (Springer, Berlin, 1988).
79. E. Mansfield, *Differential Groebner bases* (Ph D Thesis, University of Sydney, 1991).
80. V. B. Matveev and M. A. Salle, *Darboux transformations and solitons* (Springer-Verlag, Berlin, 1991).
81. D. W. McLaughlin and A. C. Scott, A restricted Bäcklund transformation, *J. Math. Phys.* **14** (1973) 1817–1828.
82. H. C. Morris, Prolongation structures and a generalized inverse scattering problem, *J. Math. Phys.* **17** (1976) 1867–1869.
83. M. Musette, Insertion of the Darboux transformation in the invariant Painlevé analysis of nonlinear partial differential equations, *Painlevé transcendents, their asymptotics and physical applications*, 197–209, eds. D. Levi and P. Winternitz (Plenum Publishing Corp., New York, 1992).
84. M. Musette, Painlevé analysis for nonlinear partial differential equations, *The Painlevé property, one century later*, 517–572, ed. R. Conte, CRM series in mathematical physics (Springer, New York, 1999) solv-int/9804003.
85. M. Musette, Chapter ?, this volume.
86. M. Musette and R. Conte, Algorithmic method for deriving Lax pairs from the invariant Painlevé analysis of nonlinear partial differential equations, *J. Math. Phys.* **32** (1991) 1450–1457.
87. M. Musette and R. Conte, Solitary waves and Lax pairs from polynomial expansions of nonlinear differential equations, *Nonlinear evolution equations and dynamical systems*, pp. 161–170, eds. M. Boiti, L. Martina and F. Pempinelli (World Scientific, Singapore, 1992).
88. M. Musette and R. Conte, The two-singular manifold method, I. Modified KdV and sine-Gordon equations, *J. Phys. A* **27** (1994) 3895–3913.
89. M. Musette and R. Conte, Non-Fuchsian extension to the Painlevé test, *Phys. Lett. A* **206** (1995) 340–346.
90. M. Musette and R. Conte, Bäcklund transformation of partial differential equations obtained from the Painlevé-Gambier classification, I. Kaup-Kupershmidt equation, *J. Math. Phys.* **39** (1998) 5617–5630.
91. M. Musette, R. Conte, and C. Verhoeven, Bäcklund transformation and nonlinear superposition formula of the Kaup-Kupershmidt and Tzitzéica equations, 19 pages, *Bäcklund and Darboux transformations : the geometry of soliton theory*, eds. C. Rogers, D. Levi, A. Coley, R. Milson, and P. Winternitz, CRM Proceedings and Lecture Notes ? (2000). American Mathematical Society, Providence, R.I. AARMS-CRM workshop (Halifax, 5–9 June 1999). S99/071.
92. A. Newell, M. Tabor, and Zeng Y.-b., A unified approach to Painlevé expansions, *Physica D* **29** (1987) 1–68.
93. A. C. Newell and J. A. Whitehead, Finite bandwidth, finite amplitude convection, *J. Fluid Mech.* **38** (1969) 279–303.
94. K. Nozaki, Hirota’s method and the singular manifold expansion, *J. Phys. Soc. Japan* **56** (1987) 3052–3054.
95. K. Nozaki and N. Bekki, Exact solutions of the generalized Ginzburg-Landau equation, *J. Phys. Soc. Japan* **53** (1984) 1581–1582.
96. M. C. Nucci and P. A. Clarkson, The nonclassical method is more general than the direct method for symmetry reductions. An example of the Fitzhugh-Nagumo equation *Phys. Lett. A* **164** (1992) 49–56.

97. P. Painlevé, Sur les équations différentielles du second ordre et d'ordre supérieur dont l'intégrale générale est uniforme, *Acta Math.* **25** (1902) 1–85.
98. P. Painlevé, Sur les équations différentielles du second ordre à points critiques fixes, *C. R. Acad. Sc. Paris* **143** (1906) 1111–1117.
99. N. R. Pereira and L. Stenflo, Nonlinear Schrödinger equation including growth and damping, *Phys. Fluids* **20** (1977) 1733–1743.
100. A. Pickering, A new truncation in Painlevé analysis, *J. Phys. A* **26** (1993) 4395–4405.
101. A. Pickering, The singular manifold method revisited, *J. Math. Phys.* **37** (1996) 1894–1927.
102. A. Ramani, B. Grammaticos, and T. Bountis, The Painlevé property and singularity analysis of integrable and nonintegrable systems, *Physics Reports* **180** (1989) 159–245.
103. C. Rogers and S. Carillo, On reciprocal properties of the Caudrey-Dodd-Gibbon and Kaup-Kupershmidt hierarchies, *Physica Scripta* **36** (1987) 865–869.
104. C. Rogers and W. F. Shadwick, *Bäcklund transformations and their applications* (Academic press, New York, 1982).
105. J. Satsuma, Exact solutions of Burgers' equation with reaction terms, *Topics in soliton theory and exact solvable nonlinear equations*, 255–262, eds. M. J. Ablowitz, B. Fuchssteiner, and M. D. Kruskal (World Scientific, Singapore, 1987).
106. J. Satsuma and D. J. Kaup, A Bäcklund transformation for a higher order Korteweg-de Vries equation, *J. Phys. Soc. Japan* **43** (1977) 692–697.
107. C. Scheen and J. Demaret, Analytic structure and chaos in Bianchi type-IX relativistic dynamics, *Phys. Lett. B* **399** (1997) 207–214.
108. W. Schief, The Tzitzéica equation: a Bäcklund transformation interpreted as truncated Painlevé expansion, *J. Phys. A* **29** (1996) 5153–5155.
109. K. M. Tamizhmani and M. Lakshmanan, Linearization and Painlevé property of Liouville and Cheng equations, *J. Math. Phys.* **27** (1986) 2257–2258.
110. G. Tzitzéica, Sur une nouvelle classe de surfaces, *Rendiconti del Circolo Matematico di Palermo* **25** (1908) 180–187.
111. G. Tzitzéica, Sur une nouvelle classe de surfaces, *C. R. Acad. Sc. Paris* **150** (1910) 955–956.
112. H. D. Wahlquist and F. B. Estabrook, Prolongation structures of nonlinear evolution equations, *J. Math. Phys.* **16** (1975) 1–7.
113. J. Weiss, The Painlevé property for partial differential equations. II: Bäcklund transformation, Lax pairs, and the Schwarzian derivative, *J. Math. Phys.* **24** (1983) 1405–1413.
114. J. Weiss, Bäcklund transformation and linearizations of the Hénon-Heiles system, *Phys. Lett. A* **102** (1984) 329–331.
115. J. Weiss, On classes of integrable systems and the Painlevé property, *J. Math. Phys.* **25** (1984) 13–24.
116. J. Weiss, The sine-Gordon equations : Complete and partial integrability, *J. Math. Phys.* **25** (1984) 2226–2235.
117. J. Weiss, The Painlevé property and Bäcklund transformations for the sequence of Boussinesq equations, *J. Math. Phys.* **26** (1985) 258–269.
118. J. Weiss, Bäcklund transformation and the Painlevé property, *J. Math. Phys.* **27** (1986) 1293–1305.

119. J. Weiss, M. Tabor and G. Carnevale, The Painlevé property for partial differential equations, *J. Math. Phys.* **24** (1983) 522–526.
120. R. Willox, private communication (1995).
121. P. Winternitz, Chapter ?, this volume.
122. Yang Huan-xiong and Li You-quan, Prolongation approach to Bäcklund transformation of Zhiber-Mikhailov-Shabat equation, *J. Math. Phys.* **37** (1996) 3491–3497.
123. V. E. Zakharov, On stochastization of one-dimensional chains of nonlinear oscillators, *Zh. Eksp. Teor. Fiz.* **65**, 219–225 (1973) [English: *Soviet Physics JETP* **38**, 108–110 (1974)].
124. V. E. Zakharov and A. B. Shabat, A scheme for integrating the nonlinear equations of mathematical physics by the method of the inverse scattering problem. I, *Funktsional'nyi Analiz i Ego Prilozheniya* **8**, 43–53 (1974) [English: *Funct. Anal. Appl.* **8**, 226–235 (1974)].
125. A. V. Zhiber and A. B. Shabat, Klein-Gordon equations with a non-trivial group, *Dokl. Akad. Nauk SSSR* **247**, 1103–1107 (1979) [English : *Sov. Phys. Dokl.* **24**, 607–609 (1979)].