



$$1. \quad \frac{\partial w}{\partial t} = a \frac{\partial}{\partial x} \left(w^m \frac{\partial w}{\partial x} \right).$$

Heat equation with a power-law nonlinearity. This equation occurs in nonlinear problems of heat and mass transfer and flows in porous media.

1°. Solutions:

$$w(x, t) = (\pm kx + k\lambda t + A)^{1/m}, \quad k = \lambda m/a,$$

$$w(x, t) = \left[\frac{m(x - A)^2}{2a(m + 2)(B - t)} \right]^{\frac{1}{m}},$$

$$w(x, t) = \left[A|t + B|^{-\frac{m}{m+2}} - \frac{m}{2a(m + 2)} \frac{(x + C)^2}{t + B} \right]^{\frac{1}{m}},$$

$$w(x, t) = \left[\frac{m(x + A)^2}{\varphi(t)} + B|x + A|^{\frac{m}{m+1}} |\varphi(t)|^{-\frac{m(2m+3)}{2(m+1)^2}} \right]^{\frac{1}{m}}, \quad \varphi(t) = C - 2a(m + 2)t,$$

where $A, B, C,$ and λ are arbitrary constants. The second solution for $B > 0$ corresponds to blow-up regimes (the solution increases without bound on a finite time interval).

2°. There are solutions of the following forms:

$$w(x, t) = (t + C)^{-1/m} F(x) \quad \text{multiplicative separable solution;}$$

$$w(x, t) = t^\lambda G(\xi), \quad \xi = xt^{-\frac{m\lambda+1}{2}} \quad \text{self-similar solution;}$$

$$w(x, t) = e^{-2\lambda t} H(\eta), \quad \eta = xe^{\lambda mt} \quad \text{generalized self-similar solution;}$$

$$w(x, t) = (t + C)^{-1/m} U(\zeta), \quad \zeta = x + \lambda \ln(t + C),$$

where $C, \beta,$ and λ are arbitrary constants.

References

- Zel'dovich, Ya. B. and Kompaneets, A. S.,** On the theory of propagation of heat with the heat conductivity depending upon the temperature, pp. 61–71, In: *Collection in Honor of the Seventieth Birthday of Academician A. F. Ioffe* [in Russian], Izdat. Akad. Nauk SSSR, Moscow, 1950.
- Ovsiannikov, L. V.,** Group properties of nonlinear heat equations [in Russian], *Doklady AN SSSR*, Vol. 125, No. 3, pp. 492–495, 1959.
- Ibragimov, N. H.** (Editor), *CRC Handbook of Lie Group Analysis of Differential Equations, Vol. 1, Symmetries, Exact Solutions and Conservation Laws*, CRC Press, Boca Raton, 1994.
- Samarskii, A. A., Galaktionov, V. A., Kurdyumov, S. P., and Mikhailov, A. P.,** *Blow-up in Problems for Quasilinear Parabolic Equations*, Walter de Gruyter, Berlin, 1995.
- Rudykh, G. A. and Semenov, E. I.,** On new exact solutions of a one-dimensional nonlinear diffusion equation with a source (sink) [in Russian], *Zhurn. Vychisl. Matem. i Matem. Fiziki*, Vol. 38, No. 6, pp. 971–977, 1998.
- Polyanin, A. D. and Zaitsev, V. F.,** *Handbook of Nonlinear Partial Differential Equations*, Chapman & Hall/CRC, Boca Raton, 2004.