



31.  $y'_x = f\left(\frac{ax + by + c}{\alpha x + \beta y + \gamma}\right).$

1°. For  $\Delta = a\beta - b\alpha \neq 0$ , the transformation

$$x = u + \frac{b\gamma - c\beta}{\Delta}, \quad y = v(u) + \frac{c\alpha - a\gamma}{\Delta}$$

leads to an equation:

$$v'_u = f\left(\frac{au + bv}{\alpha u + \beta v}\right).$$

Dividing both the numerator and denominator of the fraction on the right-hand side by  $u$ , we obtain a homogeneous equation of the form 1.5.

2°. For  $\Delta = 0$  and  $b \neq 0$ , the substitution  $v(x) = ax + by + c$  leads to a separable equation:

$$v'_x = a + bf\left(\frac{bv}{\beta v + b\gamma - c\beta}\right).$$

3°. For  $\Delta = 0$  and  $\beta \neq 0$ , the substitution  $v(x) = \alpha x + \beta y + \gamma$  also leads to a separable equation:

$$v'_x = \alpha + \beta f\left(\frac{bv + c\beta - b\gamma}{\beta v}\right).$$

## References

- Kamke, E.**, *Differentialgleichungen: Lösungsmethoden und Lösungen, I, Gewöhnliche Differentialgleichungen*, B. G. Teubner, Leipzig, 1977.
- Polyanin, A. D. and Zaitsev, V. F.**, *Handbook of Exact Solutions for Ordinary Differential Equations, 2nd Edition*, Chapman & Hall/CRC, Boca Raton, 2003.