



2. $y''_{xx} - ax^n y = 0$.

1°. Solution for $n = -2$:

$$y = C_1|x|^{k_1} + C_2|x|^{k_2},$$

where C_1 and C_2 are arbitrary constants, k_1 and k_2 are roots of the quadratic equation $k^2 - k - a = 0$.

2°. Assume $2/(n + 2) = 2m + 1$, where m is an integer. Then the solution is:

$$y = \begin{cases} x(x^{1-2q}D)^{m+1} \left[C_1 \exp\left(\frac{\sqrt{a}}{q}x^q\right) + C_2 \exp\left(-\frac{\sqrt{a}}{q}x^q\right) \right] & \text{if } m \geq 0, \\ (x^{1-2q}D)^{-m} \left[C_1 \exp\left(\frac{\sqrt{a}}{q}x^q\right) + C_2 \exp\left(-\frac{\sqrt{a}}{q}x^q\right) \right] & \text{if } m < 0, \end{cases}$$

where $D = \frac{d}{dx}$, $q = \frac{n+2}{2} = \frac{1}{2m+1}$.

3°. For any n , the solution is expressed in terms of the Bessel functions and modified Bessel functions:

$$y = \begin{cases} C_1\sqrt{x} J_{\frac{1}{2q}}\left(\frac{\sqrt{-a}}{q}x^q\right) + C_2\sqrt{x} Y_{\frac{1}{2q}}\left(\frac{\sqrt{-a}}{q}x^q\right) & \text{if } a < 0, \\ C_1\sqrt{x} I_{\frac{1}{2q}}\left(\frac{\sqrt{a}}{q}x^q\right) + C_2\sqrt{x} K_{\frac{1}{2q}}\left(\frac{\sqrt{a}}{q}x^q\right) & \text{if } a > 0, \end{cases}$$

where $q = \frac{1}{2}(n + 2)$.

References

Kamke, E., *Differentialgleichungen: Lösungsmethoden und Lösungen, I. Gewöhnliche Differentialgleichungen*, B. G. Teubner, Leipzig, 1977.
Polyanin, A. D. and Zaitsev, V. F., *Handbook of Exact Solutions for Ordinary Differential Equations, 2nd Edition*, Chapman & Hall/CRC, Boca Raton, 2003.