



12.
$$\frac{\partial u}{\partial t} = a \frac{\partial^2 u}{\partial x^2} + u f\left(\frac{w}{u}\right) - w g\left(\frac{w}{u}\right) + \frac{u}{\sqrt{u^2 + w^2}} h\left(\frac{w}{u}\right),$$
$$\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + w f\left(\frac{w}{u}\right) + u g\left(\frac{w}{u}\right) + \frac{w}{\sqrt{u^2 + w^2}} h\left(\frac{w}{u}\right).$$

Solution:

$$u = r(x, t) \cos \varphi(t), \quad w = r(x, t) \sin \varphi(t),$$

where the function $\varphi = \varphi(t)$ is determined by the separable first-order ordinary differential equation

$$\varphi'_t = g(\tan \varphi),$$

and the function $r = r(x, t)$ satisfies the linear equation

$$\frac{\partial r}{\partial t} = a \frac{\partial^2 r}{\partial x^2} + r f(\tan \varphi) + h(\tan \varphi). \tag{1}$$

The change of variable

$$r = F(t) \left[Z(x, t) + \int \frac{h(\tan \varphi) dt}{F(t)} \right], \quad F(t) = \exp \left[\int f(\tan \varphi) dt \right]$$

brings (1) to the linear heat equation

$$\frac{\partial Z}{\partial t} = a \frac{\partial^2 Z}{\partial x^2}.$$